

ON L^2_W -QUASI-DERIVATIVES FOR SOLUTIONS OF PERTURBED GENERAL QUASI-DIFFERENTIAL EQUATIONS

SOBHY EL-SAYED IBRAHIM

Abstract. This paper is concerned with square integrable quasi-derivatives for any solution of a general quasi-differential equations of n th order with complex coefficients $M[y] - \lambda \omega y = \omega f(t, y^{[0]}, \dots, y^{[n-1]})$, $t \in [a, b)$ provided that all r th quasi-derivatives of solutions of $M[y] - \lambda \omega y = 0$ and all solutions of its formal adjoint $M^+[z] - \lambda \omega z = 0$ are in $L^2_\omega(a, b)$ and under suitable conditions on the function f .

1. Introduction

In [16] Anton Zettl proved, under suitable conditions on f , that $y^{(j)} \in L^2[0, \infty)$, $j = 0, 1, \dots, n - 1$ for any solution y of $M[y] = f(t, y)$ provided that all j th derivatives of solutions of $M[y] = 0$ and all solutions of $M^+[y] = 0$ are in $L^2[0, \infty)$, when M be a regular ordinary linear differential operator of order n with coefficients which are locally integrable on $[0, \infty)$ and M^+ is the formal adjoint of M . The case $j = 0$ was considered in [16] for general n th order M and for $n = 2$ by Bradley [1]. Also, in [14] Wong proved that all solutions of a perturbed linear differential equation belong to $L^2[0, \infty)$ assuming the fact that all solutions of the unperturbed equation possess the same property, these results generalized by Ibrahim in [10] for general ordinary quasi-differential equations of n th order.

Our objective in this paper is to extend the results of Ibrahim, Wong and Zettl in [10], [14], [15] and [16] for general ordinary quasi-differential expression M of order n with complex coefficients and to prove under suitable conditions on f , that the quasi-derivatives $y^{[r]} \in L^2_\omega(a, b)$, $r = 0, 1, \dots, n - 1$ for any solution y of

$$M[y] - \lambda \omega y = \omega f(t, y^{[0]}, \dots, y^{[n-1]})(\lambda \in \mathbb{C}) \text{ on } [a, b), \quad (1.1)$$

provided that all r th quasi-derivatives of solutions of the equation

$$M[y] - \lambda \omega y = 0 \quad (1.2)$$

Received September, 18, 1997.

1991 *Mathematics Subject Classification.* Primary 34A05, 34B25, 34C11, 34E10, 34E15, 34G10, 47A55 and 47E05.

Key words and phrases. Quasi-differential operators, regular, singular, bounded and square integrable solutions.

and all solutions of

$$M^+[y] - \bar{\lambda} \omega y = 0 \tag{1.3}$$

are in $L^2_w(a, b)$. These results are extensions of those proved by Ibrahim in [10]. Our approach is based on an extension of Gronwall’s inequality used by Bradley and due to Gollwitzer [6], on a technical lemma from Goldberg’s book [5] and on an appropriate formulation of the variation of parameters formula.

We deal throughout with a quasi-differential expression M of arbitrary order n defined by a Shin-Zettl matrix on the interval $I = (a, b)$. The left-hand end-point of I is assumed to be regular but the right-hand end-point may be either regular or singular.

2. Notation and Preliminaries

The set $Z_n(I)$ of Shin-Zettl matrices on the interval I consists of $n \times n$ - matrices $A = \{a_{rs}\}$ whose entries are complex-valued functions on I which satisfy the following conditions:

$$\begin{cases} a_{rs} \in L^1_{loc}(I) & (1 \leq r, s \leq n, n \geq 2) \\ a_{r,r+1} \neq 0 \text{ a.e. on } I & (1 \leq r \leq n-1) \\ a_{rs} = 0 \text{ a.e. on } I & (2 \leq r+1 < s \leq n) \end{cases} \tag{2.1}$$

For $A \in Z_n(I)$, the quasi-derivatives associated with A are defined by

$$\begin{cases} y^{[0]} := y \\ y^{[r]} := a_{r,r+1}^{-1} \{ (y^{[r-1]})' - \sum_{s=1}^r a_{rs} y^{[s-1]} \}, & (1 \leq r \leq n-1) \\ y^{[n]} := (y^{[n-1]})' - \sum_{s=1}^n a_{ns} y^{[s-1]} \end{cases} \tag{2.2}$$

where the prime $'$ denotes differentiation.

The quasi-differential expression M associated with A is given by

$$M[y] = i^n y^{[n]}, \quad (n \geq 2) \tag{2.3}$$

this being defined on the set

$$V(M) := \{y : y^{[r-1]} \in AC_{loc}(I), \quad r = 1, 2, \dots, n\}, \tag{2.4}$$

where $AC_{loc}(I)$ denotes the set of all functions which are absolutely continuous on every compact subinterval of I .

The formal adjoint M^+ of M is defined by the matrix $A^+ \in Z_n(I)$ given by,

$$A^+ = -L^{-1}A^*L, \tag{2.5}$$

where A^* is the conjugate transpose of A and L is the non-singular $n \times n$ - matrix,

$$L := \{(-1)^r \delta_{r,n+1-s}\} \quad (1 \leq r, s \leq n), \tag{2.6}$$

δ being the kronecker delta. If $A^+ = \{a_{rs}^+\}$, then it follows that

$$a_{rs}^+ = (-1)^{r+s+1} \bar{a}_{n-s+1, n-r+1}' \quad \text{for each } r \text{ and } s. \tag{2.7}$$

The quasi-derivatives associated with A^+ are therefore.

$$\begin{cases} y_+^{[0]} := y \\ y_+^{[r]} := (\bar{a}_{n-r, n-r+1})^{-1} \{ (y_+^{[r-1]})' - \sum_{s=1}^r (-1)^{r+s+1} \bar{a}_{n-s+1, n-r+1} y_+^{[s-1]} \}, \quad (1 \leq r \leq n-1) \\ y_+^{[n]} := (y_+^{[n-1]})' - \sum_{s=1}^n (-1)^{n+s+1} \bar{a}_{n-s+1, 1} y_+^{[s-1]} \end{cases} \tag{2.8}$$

and

$$M^+[y] = i^n y_+^{[n]} \quad (n \geq 2), \tag{2.9}$$

for all y in

$$V(M^+) := \{y : y_+^{[r-1]} \in AC_{loc}(I), r = 1, 2, \dots, n\}. \tag{2.10}$$

Note that $(A^+)^+ = A$ and so $(M^+)^+ = M$. We refer to [3], [9] and [17] for a full account of the above and subsequent result on quasi-differential expressions.

Let the interval I have end-points a and b , $-\infty \leq a < b \leq \infty$, and let w be a function which satisfies,

$$w > 0 \text{ almost everywhere on } I, \quad w \in L^1_{loc}(I). \tag{2.11}$$

The equation

$$M[y] = \lambda w y, \quad (\lambda \in \mathbb{C}) \tag{2.12}$$

on I is said to be *regular* at the left end-point a if for all $x \in (a, b)$,

$$a \in \mathbb{R}, a_{rs} \in L^1[a, x], \quad (r, s = 1, 2, \dots, n). \tag{2.13}$$

Otherwise (2.12) is said to be *singular* at a . Similarly we define the terms regular and singular at b . If (2.12) is regular at both end-points, then it is said to be regular; in this case we have

$$a, b \in \mathbb{R}, a_{rs} \in L^1(a, b), \quad (r, s = 1, 2, \dots, n); \tag{2.14}$$

see [2], [8] and [9].

We shall be concerned with the case when a is a regular end-point for (2.12), but the end-point b being allowed to be either regular or singular. Note that in view of (2.7), an end-point of the interval I is regular for (2.12) if and only if it is regular for the equation,

$$M^+[y] = \bar{\lambda} w y \quad (\lambda \in \mathbb{C}) \tag{2.15}$$

Let $L^2_w(a, b)$ denote the usual weighted L^2 -space with inner-product

$$(f, g) = \int_a^b f(x) \overline{g(x)} w(x) dx, \tag{2.16}$$

and norm $\|f\| := (f, f)^{1/2}$, this is a Hilbert space on identifying functions which differ only on null space.

Denote by $S(M)$ and $S(M^+)$ the sets of all solutions of $M[y] - \lambda\omega y = 0$ and $M^+[y] - \bar{\lambda}\omega y = 0$, respectively, and $S^r(M) = \{y^{[r]} | M[y] - \lambda\omega y = 0, r = 0, 1, \dots, n - 1\}$ denotes the set of all quasi-derivatives of solutions of $M[y] - \lambda\omega y = 0$, etc. Let $\phi_k(t, \lambda)$ for $k = 1, 2, \dots, n$ be the solutions of the homogeneous equation (1.2) determined by the initial conditions,

$$\phi_k^{[r]}(t_0, \lambda) = \delta_{k,r+1} \quad \text{for all } t_0 \in [a, b), \tag{2.17}$$

($k = 1, 2, \dots, n; r = 0, 1, \dots, n - 1$). Then $\phi_k^{[r]}(t_0, \lambda)$ is continuous in (t, λ) for $a < t < b$, $|\lambda| < \infty$, and for fixed t it is entire in λ . Let $\phi_k^+(t, \lambda)$ for $k = 1, 2, \dots, n$ be the solutions of the homogeneous equation (1.3) determined by the initial conditions,

$$(\phi_k^+)^{[r]}(t_0) = (-1)^{k+r} \delta_{k,n-r'}, \quad \text{for all } t_0 \in [a, b), \tag{2.18}$$

($k = 1, 2, \dots, n; r = 0, 1, \dots, n - 1$).

Suppose $a < t_0 < b$. According to Gilbert [4, Section 3] and Ibrahim [11, Section 3], a solution of $M[\phi] - \lambda\omega\phi = \omega f$, $f \in L^1_\omega(a, b)$ satisfying $\phi^{[r]}(t_0, \lambda) = \alpha_{r+1}$, $r = 0, 1, \dots, n - 1$, is given by

$$\begin{aligned} \phi(t, \lambda) = & \sum_{j=1}^n \alpha_j(\lambda) \phi_j(t, \lambda_0) + [(\lambda - \lambda_0)/(i^n)] \sum_{j,k=1}^n \xi^{jk} \phi_j(t, \lambda_0) \\ & \cdot \int_{t_0}^t \phi_j^+(s, \lambda_0) f(s) \omega(s) ds \end{aligned} \tag{2.19}$$

where $\phi_k^+(t, \lambda)$ stands for the complex conjugate of $\phi_k(t, \lambda)$, and for each j, k , ξ^{jk} is a constant which is independent of t, λ (but does depend in general on t), also see [9, Corollary 3.10] and [17, Theorem 3].

Theorem 2.1. (Existence and Uniqueness Theorem). *Suppose $f \in L^1_\omega(a, b)$ and suppose that the conditions (2.1) are satisfied. Then given any complex numbers $\alpha_j \in \mathbb{C}, j = 0, \dots, n - 1$ and $t_0 \in (a, b)$ there exists a unique solution of $M[\phi] - \lambda\phi\omega = \omega f$ in (a, b) which satisfies*

$$\phi^{[r]}(t_0, \lambda) = \alpha_{r+1}, \quad r = 0, \dots, n - 1.$$

Proof. See [2, Theorem 3.10.1], [8, Theorem 16.2.2] and [9, Theorem 1.11].

3. Some Technical Lemmas

Our first lemma is a form of the variation of parameters formula. However it is different from the form of this formula generally found in textbooks and the literature. For the variation of parameters formula for general quasi-differential equations, see [11, Section 3] and [17, Theorem 3]. These contain Lemma 3.1 as a special case, see [16].

Lemma 3.1. *Suppose f is locally $L^1_w(a, b)$ function and $\phi(t, \lambda)$ is the solution of $M[y] - \lambda wy = wf$ satisfying*

$$\phi^{[r]}(t_0, \lambda) = \alpha_{r+1}(\lambda), \quad t_0 \in [a, b] \quad \text{for all } r = 0, 1, \dots, n - 1.$$

Then,

$$\begin{aligned} \phi^{[r]}(t, \lambda) = & \sum_{j=1}^n \alpha_j(\lambda) \phi_j^{[r]}(t, \lambda_0) + \left[(\lambda - \lambda_0) / (i^n) \right] \sum_{j,k=1}^n \xi^{jk} \phi_j^{[r]}(t, \lambda_0) \\ & \cdot \int_a^t \overline{\phi_j^+(s, \lambda_0)} f(s) w(s) ds \quad \text{for } r = 0, 1, \dots, n - 1. \end{aligned} \tag{3.1}$$

Crucial in the study of boundedness of solutions of quasi-differential equations is the fundamental inequality of Gronwall, see [7]. Here, we shall also need the following variant which may be found in [13].

Lemma 3.2. *Let $u(t), v(t)$ be two non-negative functions, locally integrable on $[a, b)$. Then, the following inequality for $0 \leq p < 1$,*

$$u(t) \leq C_0 + \int_a^t v(s) u^p(s) ds, \quad C_0 > 0,$$

implies that,

$$u(t) \leq \left[C_0^{1-p} + (1-p) \int_a^t v(s) ds \right]^{1/(1-p)} \tag{3.2}$$

In particular, if $v(t) \in L^1(a, b)$, then (3.2) implies that $u(t)$ is bounded.

The next lemma is a special case of an extension of the well-known Gronwall inequality due to Gollwitzer [6] (See also Willett [12] and Willet-Wong [13]).

Lemma 3.3. *Let u, z, g, h , be continuous non-negative functions on $[a, b)$ and suppose that*

$$u(t) \leq z(t) + g(t) \left[\int_a^t u^2(s) h(s) ds \right]^{\frac{1}{2}} \quad \text{for } t \geq a.$$

Then,

$$u(t) \leq z(t) + g(t) \left[\int_a^t 2z^2(s) h(s) \exp \left[\int_a^s 2g^2(x) h(x) dx \right] ds \right]^{\frac{1}{2}} \quad \text{for } t \geq a.$$

The next two Lemmas has been proved in [9] for general case.

Lemma 3.4. ([9, Proposition 3.23]) *If all solutions of $M[\phi] = \lambda_0 w \phi$ are bounded on $[a, b)$ for some $\lambda_0 \in \mathbb{C}$. and $\phi_j^+ \in L^1_w(a, b)$ for $j = 1, 2, \dots, n$, then all solutions of $M[\phi] = \lambda w \phi$ are also bounded on $[a, b)$ for all $\lambda \in \mathbb{C}$.*

Lemma 3.5. ([9, Proposition 3.24]) *If all solutions of $M[\phi] = \lambda_0 w \phi$ and $M^+[\phi] = \bar{\lambda}_0 w \phi$ are in $L^2_w(a, b)$ for some $\lambda_0 \in \mathbb{C}$, then all solutions of $M[\phi] = \lambda w \phi$ and $M^+[\phi] = \bar{\lambda} w \phi$ are in $L^2_w(a, b)$ for all $\lambda \in \mathbb{C}$.*

4. The Main Results

Suppose there exist non-negative continuous functions $k(t)$ and $h_i(t)$, $i = 0, \dots, n - 1$ such that

$$|f(t, y^{[0]}, \dots, y^{[n-1]})| \leq k(t) + \sum_{i=0}^{n-1} h_i(t) |y^{[i]}|^\sigma, \tag{4.1}$$

for $t \geq a$, $-\infty < y^{[i]} < \infty$, for each $i = 0, \dots, n - 1$; $0 \leq \sigma \leq 1$.

Theorem 4.1. *Suppose f satisfies (4.1) with $\sigma = 1$, $S^r(M) \cup S(M^+) \subset L^\infty(a, b)$ for some $r = 0, 1, \dots, n - 1$ and some $\lambda_0 \in \mathbb{C}$ and that,*

- (i) $k(t) \in L^1_w(a, b)$ for all $t \in [a, b)$,
- (ii) $h_i(t) \in L^1_w(a, b)$ for all $t \in [a, b)$, $i = 0, 1, \dots, n - 1$.

Then $\phi^{[r]}(t, \lambda)$ is bounded on $[a, b)$ for any solution $\phi(t, \lambda)$ of the equation (1.1) for all $\lambda \in \mathbb{C}$.

Proof. Note that (4.1) and Theorem 2.1 implies that all solutions exist on the entire interval $[a, b)$, see [2, Chapter 3], [10], [14] and [15].

Let $\{\phi_1(t, \lambda_0), \dots, \phi_n(t, \lambda_0)\}$ and $\{\phi_1^+(t, \lambda_0), \dots, \phi_n^+(t, \lambda_0)\}$ be two sets of linearly independent solutions of (1.2) and (1.3) respectively and let $\phi(t, \lambda)$ be any solution of (1.1) on $[a, b)$, then by Lemma 3.1, we have,

$$\begin{aligned} \phi^{[r]}(t, \lambda) &= \sum_{j=1}^n \alpha_j(\lambda) \phi_j^{[r]}(t, \lambda_0) + ((\lambda - \lambda_0)/(i^n)) \sum_{j,k=1}^n \xi^{jk} \phi_j^{[r]}(t, \lambda_0) \\ &\quad \cdot \int_a^t \overline{\phi_j^+(s, \lambda_0)} f(s, \phi^{[0]}, \dots, \phi^{[n-1]}) w(s) ds, \quad \text{for } r = 0, \dots, n - 1. \end{aligned} \tag{4.2}$$

Hence,

$$\begin{aligned} |\phi^{[r]}(t, \lambda)| &\leq \sum_{j=1}^n |\alpha_j(\lambda)| |\phi_j^{[r]}(t, \lambda_0)| + |\lambda - \lambda_0| \sum_{j,k=1}^n |\xi^{jk}| |\phi_j^{[r]}(t, \lambda_0)| \\ &\quad \cdot \int_a^t \overline{|\phi_k^+(s, \lambda_0)|} \left(k(s) + \sum_{i=0}^{n-1} h_i(s) |\phi^{[i]}(s, \lambda)| \right) w(s) ds, \quad r = 0, 1, \dots, n - 1. \end{aligned} \tag{4.3}$$

Since, $k(t) \in L^1_w(a, b)$ and $\phi_j^+(t, \lambda_0)$ bounded on $[a, b)$ for some $\lambda_0 \in \mathbb{C}$, then $\phi_j^+ k \in L^1_w(a, b)$, $j = 1, 2, \dots, n$ for some $\lambda_0 \in \mathbb{C}$.

Setting,

$$C_j = |\lambda - \lambda_0| \sum_{j,k=1}^n |\xi^{jk}| \int_a^b \overline{|\phi_j^+(s, \lambda_0)|} k(s) w(s) ds, \quad (j = 1, 2, \dots, n), \tag{4.4}$$

then,

$$|\phi^{[r]}(t, \lambda)| \leq \sum_{j=1}^n (c_j + |\alpha_j(\lambda)|) |\phi_j^{[r]}(t, \lambda_0)| + |\lambda - \lambda_0| \sum_{j,k=1}^n \sum_{i=0}^{n-1} |\xi^{jk}|$$

$$|\phi_j^{[r]}(t, \lambda_0)| \int_a^t \overline{|\phi_j^+(s, \lambda_0)|} h_i(s) |\phi^{[i]}(s, \lambda)| w(s) ds, \quad r = 0, 1, \dots, n - 1. \quad (4.5)$$

By hypothesis, there exist positive constants K_0 and K_1 such that,

$$|\phi_j^{[r]}(t, \lambda_0)| \leq K_0 \quad \text{and} \quad |\phi_j^+(t, \lambda_0)| \leq K_1, \quad \text{for all } t \in [a, b],$$

$j = 1, 2, \dots, n$ and some $r = 0, 1, \dots, n - 1$.

Hence, by summing both sides of (4.5) from $r = 0, \dots, n - 1$ we get,

$$\begin{aligned} \left(\sum_{r=0}^{n-1} |\phi^{[r]}(t, \lambda)| \right) &\leq (n - 1)K_0 \sum_{j=1}^n (C_j + |\alpha_j(\lambda)|) + (n - 1)K_0K_1|\lambda - \lambda_0| \\ &\cdot \sum_{j,k=1}^n |\xi^{jk}| \int_a^t \left(\max_{0 \leq i \leq n-1} h_i(s) \right) \left(\sum_{i=0}^{n-1} |\phi^{[i]}(s, \lambda)| \right) w(s) ds. \end{aligned} \quad (4.6)$$

On application of Gronwall's inequality to (4.6) and from (ii), we deduce that $\sum_{r=0}^{n-1} |\phi^{[r]}(t, \lambda)|$ is finite and hence the result.

Remark. From [16, Section 3] and [9, Lemma 3.3], ϕ and $\phi^{[j]} \in L^2_w(a, b)$ implies that $\phi^{[r]} \in L^2_w(a, b)$ for any solution ϕ of the equation (1.1), for all $r = 1, \dots, j - 1, 1 \leq j \leq n - 1$.

Theorem 4.2. Suppose f satisfies (4.1) with $\sigma = 1, S^r(M) \cup S(M^+) \subset L^2_w(a, b)$ for some $\lambda_0 \in \mathbb{C}$ and some $r = 0, 1, \dots, n - 1$ and that (i) $k(t) \in L^2_w(a, b)$ (ii) $h_i(t) \in L^\infty(a, b), i = 0, 1, \dots, n - 1$, for all $t \in [a, b)$. Then $\phi^{[r]}(t, \lambda) \in L^2_w(a, b)$ for any solution $\phi(t, \lambda)$ of the equation (1.1) for all $\lambda \in \mathbb{C}$.

Proof. On application of the Cauchy Schwartz inequality to the integral in (4.5) we get,

$$\begin{aligned} |\phi^{[r]}(t, \lambda)| &\leq \sum_{j=1}^n (C_j + |\alpha_j(\lambda)|) |\phi_j^{[r]}(t, \lambda_0)| + |\lambda - \lambda_0| \sum_{j,k=1}^n \sum_{i=0}^{n-1} |\xi^{jk}| \\ &\cdot |\phi_j^{[r]}(t, \lambda_0)| \left(\int_a^t \overline{|\phi_j^+(s, \lambda_0)|}^2 |h_i(s)| w ds \right)^{\frac{1}{2}} \left(\int_a^t |h_i(s)| |\phi^{[i]}(s, \lambda)|^2 w ds \right)^{\frac{1}{2}}, \end{aligned} \quad (4.7)$$

$r = 0, 1, \dots, n - 1$.

Since $\phi_j^+(t, \lambda_0) \in L^2_w(a, b), j = 1, 2, \dots, n$ for some $\lambda_0 \in \mathbb{C}$ and $h_i(t) \in L^\infty(a, b)$ by hypothesis, then $\phi_j^+(t, \lambda_0) |h_i(t)|^{1/2} \in L^2_w(a, b), j = 1, 2, \dots, n, i = 0, 1, \dots, n - 1$. Let,

$$D_{ji} = \left(\int_a^t \overline{|\phi_j^+(s, \lambda_0)|}^2 |h_i(s)| w(s) ds \right)^{\frac{1}{2}}, \quad z(t) = \sum_{j=1}^n (C_j + |\alpha_j(\lambda)|) |\phi_j^{[r]}(t, \lambda_0)|,$$

and $G(t) = |\lambda - \lambda_0| \sum_{j,k=1}^n \sum_{i=0}^{n-1} D_{ji} |\xi^{jk}| |\phi_j^{[r]}(t, \lambda_0)|.$

From Lemma 3.3 we have,

$$|\phi^{[r]}(t, \lambda)| \leq Z(t) + G(t) \left(\int_a^t 2Z^2(s)|h_i(s)| \exp\left[\int_a^s 2G^2(x)|h_i(x)|w dx\right] w ds \right)^{\frac{1}{2}}.$$

Since $\int_a^b Z^2(s)|h_i(s)|w(s)ds$ and $\int_a^b G^2(x)|h_i(x)|w(x)dx$ are both finite, we conclude that $|\phi^{[r]}(t, \lambda)|$ is bounded by a linear combination of the $L_w^2(a, b)$ functions $Z(t)$ and $G(t)$. Therefore, by using Lemma 3.5, $\phi^{[r]}(t, \lambda) \in L_w^2(a, b)$ for all $\lambda \in \mathbb{C}$.

Remark. If we use the Cauchy Schwartz inequality to the integral in (4.5) as:

$$\int_a^t \overline{|\phi_j^+|} |h_i| |\phi| w ds \leq \left(\int_a^t \overline{|\phi_j^+|^2} |h_i|^2 w ds \right)^{\frac{1}{2}} \left(\int_a^t |\phi^{[i]}|^2 w ds \right)^{\frac{1}{2}}, i = 0, \dots, n - 1,$$

we get also the result. We refer to [14] and [15] for more details.

Corollary 4.3. Suppose that $f(t, y^{[0]}, \dots, y^{[n-1]}) = \sum_{i=1}^{n-1} h_i(t)y^{[i]}$, $S^r(M) \cup S(M^+) \subset L_w^2(a, b)$ for some $\lambda_0 \in \mathbb{C}$ and some $r = 0, 1, \dots, n - 1$ and that $h_i(t) \in L_w^p(a, b)$ for some $p \geq 2$, $t \in [a, b]$; $i = 0, 1, \dots, n - 1$. Then $\phi^{[r]}(t, \lambda) \in L_w^1(a, b)$ for any solution $\phi(t, \lambda)$ of the equation (1.1) for all $\lambda \in \mathbb{C}$ and all $r = 0, 1, \dots, n - 1$.

Proof. The proof is similar to Theorem 4.2 and therefore omitted.

The special case $h_i(t) \equiv 0$, $i = 0, \dots, n - 1$ and $k \in L_w^2(a, b)$ yields:

Corollary 4.4. If all solutions of $M[\phi] = \lambda_0 w \phi$ and $M^+[\phi] = \bar{\lambda}_0 w \phi$ are in $L_w^2(a, b)$ for some $\lambda_0 \in \mathbb{C}$ and $k \in L_w^2(a, b)$, then all solutions of $M[\phi] - \lambda w \phi = w k$ are in $L_w^2(a, b)$ for all $\lambda \in \mathbb{C}$.

Next, we consider (4.1) with $0 \leq \sigma < 1$, and have the following:

Theorem 4.5. Suppose f satisfies (4.1) with $0 \leq \sigma < 1$, $S^r(M) \cup S(M^+) \subset L_w^2(a, b)$ for some $\lambda_0 \in \mathbb{C}$ and some $r = 0, 1, \dots, n - 1$ and that,

- (i) $k(t) \in L_w^2(a, b)$ for all $t \in [a, b]$,
- (ii) $h_i(t) \in L_w^{2/(1-\sigma)}(a, b)$ $0 \leq \sigma < 1$, $i = 0, 1, \dots, n - 1$.

Then $\phi^{[r]}(t, \lambda) \in L_w^2(a, b)$ for any solution $\phi(t, \lambda)$ of the equation (1.1) for all $\lambda \in \mathbb{C}$.

Proof. For $0 \leq \sigma < 1$, the proof is the same up to (4.5). In this case (4.5) becomes,

$$|\phi^{[r]}(t, \lambda)| \leq \sum_{j=1}^n (c_j + |\alpha_j(\lambda)|) |\phi_j^{[r]}(t, \lambda_0)| + |\lambda - \lambda_0| \sum_{j,k=1}^n \sum_{i=0}^{n-1} |\xi^{jk}| \cdot |\phi_j^{[r]}(t, \lambda_0)| \int_a^t \overline{|\phi_j^+(s, \lambda_0)|} |h_i(s)| |\phi^{[i]}(s, \lambda)|^\sigma w(s) ds, r = 0, 1, \dots, n - 1. (4.8)$$

Applying Cauchy Schwartz inequality to the integral in (4.8), we find

$$\begin{aligned} & \int_a^t \overline{|\phi_j^+(s, \lambda_0)|} |h_i(s)| |\phi(s, \lambda)|^\sigma w(s) ds \\ & \leq \left(\int_a^t \overline{|\phi_j^+(s, \lambda_0)|} |h_i(s)|^\mu w(s) ds \right)^{\frac{1}{\mu}} \left(\int_a^t |\phi^{[i]}(s, \lambda)|^2 w(s) ds \right)^{\frac{\sigma}{2}}, \end{aligned} \tag{4.9}$$

where $\mu = 2/(2 - \sigma)$. Since $\phi_j^+(s, \lambda_0) \in L^2_w(a, b)$ for some $\lambda_0 \in \mathbb{C}$, $j = 1, 2, \dots, n$ and $h_i(s) \in L^{2/(1-\sigma)}_w(a, b)$ by hypothesis, then we have $\phi_j^+(s, \lambda_0)h_i(s) \in L^\mu_w(a, b)$ for some $\lambda_0 \in \mathbb{C}$, $j = 1, 2, \dots, n$; $i = 0, 1, \dots, n - 1$. Using this fact and (4.9) in (4.8), we obtain

$$|\phi^{[r]}(t, \lambda)| \leq \sum_{j=1}^n (c_j + |\alpha_j(\lambda)|) |\phi_j^{[r]}(t, \lambda_0)| + K_0 |\lambda - \lambda_0| \sum_{j,k=1}^n \sum_{i=0}^{n-1} |\xi^{jk}| \cdot |\phi_j^{[r]}(t, \lambda_0)| \left(\int_a^t |\phi^{[i]}(s, \lambda)|^2 w(s) ds \right)^{\frac{\sigma}{2}}, \quad r = 0, 1, \dots, n - 1, \quad (4.10)$$

where $K_0 = \|\phi_j^+(t, \lambda_0)h_i(s)\|_\mu$, $\|\cdot\|_\mu$ denotes the norm in $L^\mu_w(a, b)$. From the inequality

$$(u + v)^2 \leq 2(u^2 + v^2), \quad (4.11)$$

it follows that

$$|\phi^{[r]}(t, \lambda)|^2 \leq 4 \sum_{j=1}^n (c_j^2 + |\alpha_j(\lambda)|^2) |\phi_j^{[r]}(t, \lambda_0)|^2 + 4K_0^2 |\lambda - \lambda_0|^2 \sum_{j,k=1}^n \sum_{i=0}^{n-1} |\xi^{jk}|^2 \cdot |\phi_j^{[r]}(t, \lambda_0)|^2 \left(\int_a^t |\phi^{[i]}(s, \lambda)|^2 w(s) ds \right)^\sigma, \quad r = 0, 1, \dots, n - 1. \quad (4.12)$$

Setting, $K_1 = \int_a^b |\phi_j^{[r]}(t, \lambda_0)|^2 w(s) ds$, for some $\lambda_0 \in \mathbb{C}$ and some $r = 0, \dots, n - 1$; $j = 1, 2, \dots, n$ and integrate (4.12) to obtain

$$\int_a^t |\phi^{[r]}(s, \lambda)|^2 w(s) ds \leq K_2 + 4K_0^2 |\lambda - \lambda_0|^2 \left(\sum_{j,k=1}^n \sum_{i=0}^{n-1} |\xi^{jk}|^2 \int_a^t |\phi_j^{[r]}(s, \lambda_0)|^2 \cdot \left[\left(\int_a^s |\phi^{[i]}(x, \lambda)|^2 w(x) dx \right)^\sigma \right] w(s) ds, \quad (4.13)$$

where

$$K_2 = 4 \sum_{j=1}^n (c_j^2 + |\alpha_j(\lambda)|^2) K_1$$

On application of Lemma 3.2 to (4.13) for $0 \leq \sigma < 1$ and of Gronwall's inequality to (4.13) for $\sigma = 1$, yields the result.

Theorem 4.6. *Suppose f satisfies (4.1) with $0 \leq \sigma < 1$, $S^r(M) \cup S(M^+) \subset L^2_w(a, b) \cap L^\infty(a, b)$ for some $\lambda_0 \in \mathbb{C}$ and some $r = 0, 1, \dots, n - 1$ and that,*

(i) $k(t) \in L^2_w(a, b)$ for all $t \in [a, b)$,

(ii) $h_i(t) \in L^p_w(a, b)$ for any p , $1 \leq p \leq 2/(1 - \sigma)$, $i = 0, 1, \dots, n - 1$.

Then $\phi^{[r]}(t, \lambda) \in L^2_w(a, b) \cap L^\infty(a, b)$ for any solution $\phi(t, \lambda)$ of the equation (1.1) for all $\lambda \in \mathbb{C}$.

Proof. Since $S^r(M) \cup S(M^+) \subset L_w^2(a, b) \cap L^\infty(a, b)$ for some $\lambda_0 \in \mathbb{C}$ and some $r = 0, 1, \dots, n - 1$, then $\phi_j^{[r]}(t, \lambda_0), \phi_j^+(t, \lambda_0) \in L_w^q(a, b), j = 1, 2, \dots, n$ for every $q \geq 2$ and for some $\lambda_0 \in \mathbb{C}, r = 0, 1, \dots, n - 1$.

First, suppose that $h_i(t) \in L_w^p(a, b)$ for some $p, 1 \leq p \leq 2$. Setting,

$$K_0 = \|\phi^{[r]}(t, \lambda_0)\|_\infty \quad \text{and} \quad K_1 = \|\phi_j^+(t, \lambda_0)\|_\infty, j = 1, 2, \dots, n,$$

for some $\lambda_0 \in \mathbb{C}$ and some $r = 0, 1, \dots, n - 1$, we have from (4.8),

$$|\phi^{[r]}(t, \lambda)| \leq K_0 \left(\sum_{j=1}^n (c_j + |\alpha_j(\lambda)|) \right) + K_0 K_1 |\lambda - \lambda_0| \left(\sum_{j,k=1}^n \sum_{i=0}^{n-1} |\xi^{jk}| \cdot \int_a^t h_i(s) |\phi^{[i]}(s, \lambda)|^\sigma w(s) ds \right). \tag{4.14}$$

Since $h_i(t) \in L_w^p(a, b), 1 \leq p \leq 2$, then Lemma 3.2 together with Gronwall's inequality implies that $\phi^{[r]}(t, \lambda) \in L^\infty(a, b)$ for all $\lambda \in \mathbb{C}$, i.e., there exist a positive constant K_2 such that,

$$|\phi^{[r]}(t, \lambda)| \leq K_2 \text{ for all } \lambda \in \mathbb{C}, t \in [a, b), r = 0, 1, \dots, n - 1. \tag{4.15}$$

From (4.8) and (4.15), we obtain

$$|\phi^{[r]}(t, \lambda)| \leq \sum_{j=1}^n (c_j + |\alpha_j(\lambda)| + K_3) |\phi_j^{[r]}(t, \lambda_0)|,$$

for some appropriate constant K_3 . Since $\phi_j^{[r]}(t, \lambda_0) \in L_w^2(a, b)$ for some $\lambda_0 \in \mathbb{C}$ and some $r = 0, 1, \dots, n - 1$, this proves $\phi^{[r]}(t, \lambda) \in L_w^p(a, b)$ for all $\lambda \in \mathbb{C}, 1 \leq p \leq 2$.

Next, suppose that $h_i(t) \in L_w^p(a, b), 2 < p \leq 2/(1 - \sigma); i = 0, 1, \dots, n - 1$. Define $q \geq 2$ by

$$\frac{1}{q} = \frac{2 - \sigma}{2} - \frac{1}{p}$$

(which is possible because of the restriction on p). Thus $\phi_j^{[r]}(t, \lambda_0) \phi_j^+(t, \lambda_0) \in L_w^q(a, b)$ and $\phi_j^+(t, \lambda_0) h_i(t) \in L_w^\mu(a, b), \mu = 2/(2 - \sigma)$.

Repeating the same argument in the proof of Theorem 4.5, from (4.8) to (4.13), we obtain the fact that $\phi^{[r]}(t, \lambda) \in L_w^2(a, b)$. Returning to (4.9), we find that the integral on the left-hand side is bounded which implies, by (4.8) that,

$$|\phi^{[r]}(t, \lambda)| \leq \sum_{j=1}^n (c_j + |\alpha_j(\lambda)| + K_3) |\phi_j^{[r]}(t, \lambda_0)|,$$

for some appropriate constant K_3 . Since $\phi_j^{[r]}(t, \lambda_0) \in L^\infty(a, b)$, this completes the proof. We refer to [10], [14] and [16] for more details.

References

- [1] J. S. Bradley, "Comparison theorems for the square integrability of solutions of $(r(t)y')' + q(t)y = f(t, y)$," *Glasgow Math. Soc.*, 13 (1972), 75-79.
- [2] D. E. Edmunds and W. D. Evans, *Spectral Theory and Differential Operators*, Oxford University press, 1987.
- [3] W. N. Everitt and D. Race, "Some remarks on linear ordinary quasi-differential expressions," *Proc. London Math. Soc.*, (3), 54(1987), 300-320.
- [4] R. C. Gilbert, "Simplicity of linear ordinary differential operators," *Journal of Differential Equations*, 11(1972), 672-681.
- [5] S. Goldberg, "Unbounded linear operators," McGraw. Hill, New York, 1966.
- [6] H. E. Gollwitzer, "A note on a functional inequality," *Proc. Amer. Math. Soc.*, 23(1969), 642-647.
- [7] M. R. Mohana Rao, "Ordinary differential equations," *Theory and Applications*, First published in the United Kingdom in 1989 by Edward Arnold (Publishers) Limited, London.
- [8] M. N. Naimark, "Linear differential operators," G. I. T. T. L., Moscow (1954), Ungar, New York, I(1967), II(1968).
- [9] Sobhy El-sayed Ibrahim, "Problems associated with differential operators," Ph. D. thesis (1989), *Faculty of Science*, Department of Mathematics, Benha University, Egypt.
- [10] ——— "Boundedness for solutions of general ordinary quasi-differential equations," *Journal of Egyptian Mathematical Society*, 2(1994), 33-44.
- [11] ——— "The spectra of well-posed operators," *Proc. Royal Soc. of Edinburgh*, 124A(1995), 1331-1348.
- [12] D. Willett, "Nonlinear vector integral equations as contraction mappings," *Arch. Rational Mech. Anal.*, 15(1964), 79-86.
- [13] D. Willett and J. S. W. Wong, "On the discrete analogues of some generalizations of Gronwall's inequality," *Monatsh. Math.*, 69(1965), 362-367, MR 32≠2644.
- [14] J. S. W. Wong, "Square integrable solutions of perturbed linear differential equations," *Proc. Royal Society of Edinburgy*, 73A, 16, (1974/75), 251-254.
- [15] A. Zettl, "Square integrable solutions of $Ly = f(t, y)$," *Proceedings of the American Mathematical Society*, 26(1970), 635-639.
- [16] ——— "Perturbation of the limit circle case," *Quart. J. Math.*, Oxford (3), 26(1975), 355-360.
- [17] A. Zettl, "Formally self-adjoint quasi-differential operators," *Rocky Mountain Journal of Mathematics*, 5(3)(1975), 453-474.

Benha University, Faculty of Science, Department of Mathematics, Benha 13518, Egypt.