



# A novel iterative algorithm for solving variational inequality, finite family of monotone inclusion and fixed point problems

Anjali, Seema Mehra, Renu Chugh and Charu Batra

**Abstract.** In this paper, we introduce a method for finding common solution of variational inequality, finite family of monotone inclusion and fixed point problems of demicontractive mappings in a real Hilbert space. We prove strong convergence result of proposed method. We also provide a numerical example to show that our method is efficient from the numerical point of view.

**Keywords.** Iterative method, fixed point problem, variational inequality problem, monotone inclusion problem.

## 1 Introduction

Monotone inclusion problem (MIP) plays a crucial role in nonlinear analysis and optimization. MIP is the problem of finding a point  $\zeta$  in a Hilbert space  $H$  such that

$$0 \in T\zeta, \quad (1.1)$$

where  $T : H \rightarrow 2^H$  is a monotone operator. Mathematically, monotone inclusion problem includes image processing problem, variational inequality problem, split feasibility problem, convex minimization problem, equilibrium problem etc. [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13]. The first method, namely proximal point method, for solving MIP was proposed by Martinet in 1970. It was defined as

$$\zeta_{n+1} = (I + \lambda_n T)^{-1} \zeta_n. \quad (1.2)$$

Rockafeller [14] in 1976 and Bruck and Reich [15] in 1977, further developed this algorithm. But the evaluation of resolvent operator in proximal point algorithm was difficult in many cases. Consequently, to solve this issue, the operator  $T$  is divided into the sum of maximal monotone operator  $A$  and monotone operator  $B$ . As the resolvent operators  $(I + \lambda_n A)^{-1}$  and  $(I + \lambda_n B)^{-1}$  is simpler to calculate than the full resolvent  $(I + \lambda_n T)^{-1}$ . The problem (1.1) is equivalent to the following problem:

$$\text{Find } \zeta \in H \text{ such that } 0 \in (A + B)\zeta. \quad (1.3)$$

---

Received date: May 9, 2023; Published online: January 26, 2024.

2010 *Mathematics Subject Classification.* 47H05, 47H10, 47J25, 65K15.

Corresponding author: Seema Mehra.

The solution set of the problem (1.3) is given by  $(A + B)^{-1}(0)$ . The first method for solving problem (1.3) was forward-backward splitting algorithm. Many iterative algorithms have been designed to solve MIP, for instance, Douglas-Rachford splitting method [18], Peaceman-Rachford splitting method [19] and many more. Tseng [20] in 2000, introduced the modified forward-backward algorithm and proved its weak convergence. Gibali and Thong [21] in 2018, obtained a modified version of Tseng's splitting algorithm and proved its strong convergence.

Polyak [22] developed an inertial extrapolation method which is based on heavy ball method to speed up the convergence of iterative algorithms. Later, inertial extrapolation technique was used to solve MIP and numerous authors considerably enhanced it. For instance, Lorenz and Pock [23] in 2015, introduced the following inertial forward-backward algorithm for solving monotone inclusion problem and proved weak convergence in a real Hilbert space.

$$\begin{cases} \varphi_n = \zeta_n + \theta_n(\zeta_n - \zeta_{n-1}), \\ \zeta_{n+1} = (I + \lambda_n A)^{-1}(I - \lambda_n B)\varphi_n. \end{cases}$$

The above algorithm has better rate of convergence than some existing algorithms present in the literature. Thong and Chalamjiak [25] in 2019, introduced modified forward-backward splitting algorithm and proved strong convergence.

On the other hand, variational inequality is used in solving various class of problems like transportation, economics, engineering, optimization, elasticity and control theory [26, 27, 28, 29, 30]. Several numerical techniques have been devised for solving variational inequality problems (VIP). A VIP is to find a point  $\zeta$  in a convex, closed and nonempty subset  $Q$  of Hilbert space  $H$  such that

$$\langle P(\zeta), \vartheta - \zeta \rangle \geq 0 \text{ for all } \vartheta \in Q, \quad (1.4)$$

where  $P : Q \rightarrow H$  is a nonlinear mapping. The set of solutions of VIP is denoted by  $VI(Q, P)$ . Eslamian and Kamandi [31] in 2020, developed iterative algorithm for finding common solution of fixed point and monotone inclusion problem in Hilbert space. Recently, Olona and Narain [32] in 2022, introduced a method for approximating a common solution of fixed point problem for finite families of multivalued demicontractive mappings and finite families of variational inequality problem in a real Hilbert space.

Inspired and encouraged by the above results, we introduce a method for finding common solution of variational inequality problem, finite families of monotone inclusion and fixed point problems of demicontractive mappings in a real Hilbert space and prove strong convergence of proposed algorithm. Also, we provide a numerical example to show its applicability.

## 2 Preliminaries

We now give some definitions and lemmas which we will use in proving our main result. Suppose  $H$  denotes a real Hilbert space having inner product  $\langle \cdot, \cdot \rangle$  and norm  $\|\cdot\|$ . Assume that  $Q$  is a nonempty, closed and convex subset of  $H$ . In addition,  $Fix(U)$  denotes the collection of all fixed points of mapping  $U$ .

**Definition 1.** [31] “The operator  $U : H \rightarrow H$  is called

- (i) Nonexpansive, if  $\|U\zeta - U\vartheta\| \leq \|\zeta - \vartheta\|$  for all  $\zeta, \vartheta \in H$ .

- (ii) Demicontractive, if  $Fix(U) \neq \phi$  there exists  $k \in [0, 1)$  such that  $\|U\zeta - \vartheta\|^2 \leq \|\zeta - \vartheta\|^2 + k\|\zeta - U\zeta\|^2$  for all  $\zeta \in H$  and  $\vartheta \in Fix(U)$ .
- (iii) Contractive, if there exists a constant  $0 \leq \theta < 1$  such that  $\|U\zeta - U\vartheta\| \leq \theta\|\zeta - \vartheta\|$  for all  $\zeta, \vartheta \in H$ ."

**Definition 2.** [32] "Let  $Q$  be a nonempty, closed and convex subset of a real Hilbert space  $H$ . A mapping  $U : Q \rightarrow Q$  is said to be

- (i) monotone, if  $\langle U\zeta - U\vartheta, \zeta - \vartheta \rangle \geq 0$  for all  $\zeta, \vartheta \in H$ ,
- (ii)  $\alpha$ -inverse strongly monotone (ism), if there exists  $\alpha > 0$  such that  $\langle U\zeta - U\vartheta, \zeta - \vartheta \rangle \geq \alpha\|U\zeta - U\vartheta\|^2$  for all  $\zeta, \vartheta \in Q$ ,
- (iii) firmly nonexpansive, if  $\langle U\zeta - U\vartheta, \zeta - \vartheta \rangle \geq \|U\zeta - U\vartheta\|^2$  for all  $\zeta, \vartheta \in Q$ ,
- (iv) Lipschitz, if there exists a constant  $K > 0$  such that  $\|U\zeta - U\vartheta\| \leq K\|\zeta - \vartheta\|$  for all  $\zeta, \vartheta \in Q$ ."

**Definition 3.** [32] "If  $U$  is a multi-valued mapping, that is,  $U : H \rightarrow 2^H$ , then  $U$  is called monotone, if  $\langle \zeta - \vartheta, \psi - v \rangle \geq 0$  for all  $\zeta, \vartheta \in H, \psi \in U\zeta, v \in U\vartheta$  and  $U$  is maximal monotone, if the graph  $G(U)$  of  $U$  defined by  $G(U) = \{(\zeta, \vartheta) \in H \times H : \vartheta \in U(\zeta)\}$  is not properly contained in the graph of any other monotone mapping. It is generally known that  $U$  is maximal if and only if for  $(\zeta, \vartheta) \in H \times H, \langle \zeta - \psi, \vartheta - v \rangle \geq 0$  for all  $(\psi, v) \in G(U)$  implies  $\vartheta \in U(\zeta)$ ."

**Definition 4.** [32] "The metric projection  $P_Q$  is a map defined on  $H$  onto  $Q$  which assign to each  $\zeta \in H$ , the unique point in  $Q$ , denoted by  $P_Q\zeta$  such that  $\|\zeta - P_Q\zeta\| = \inf\{\|\zeta - \vartheta\| : \vartheta \in Q\}$ . It is well known that  $P_Q\zeta$  is characterized by the inequality  $\langle \zeta - P_Q(\zeta), \vartheta - P_Q(\zeta) \rangle \leq 0$ , for all  $\vartheta \in Q$  and  $P_Q$  is a firmly non-expansive mapping."

**Definition 5.** [33] "The resolvent mapping  $J_\gamma^U : H \rightarrow H$  associated with the set-value mapping  $U$  is defined by  $J_\gamma^U(\zeta) = (I + \gamma U)^{-1}(\zeta) \quad \forall \quad \zeta \in H$ , for some  $\gamma > 0$ , where  $I$  stands for the identity operator on  $H$ ."

**Definition 6.** [31] "Assume that  $T : H \rightarrow H$  is a nonlinear mapping with  $Fix(T) \neq \phi$ . Then  $I - T$  is said to be demiclosed at zero if for any  $\{\chi_n\}$  in  $H$ , the following implications holds:  $\zeta_n \rightarrow \iota$  and  $(I - T)\zeta_n \rightarrow 0 \implies \iota \in Fix(T)$ ."

**Lemma 2.1.** [34] "Assume that  $\{t_n\} \subset [0, \infty)$  is a sequence of real numbers. Suppose that

$$s_{n+1} \leq (1 - \gamma_n)s_n + \gamma_n t_n \text{ for all } n \in \mathbb{N},$$

where  $\{\gamma_n\} \subset [0, 1]$  and  $\{t_n\} \subset (-\infty, \infty)$  satisfying following conditions:

1.  $\sum_{n=1}^{\infty} \gamma_n = \infty$ ,
2.  $\limsup_{n \rightarrow \infty} t_n \leq 0$ .

Then  $\lim_{n \rightarrow \infty} s_n = 0$ ."

**Proposition 2.1.** [35] "Let  $L : Q \rightarrow H$  be an inverse strongly monotone(ism) mapping. Then,  $z \in VI(Q, L) \iff z = P_Q(z - \lambda Lz), \lambda > 0$ ."

**Proposition 2.2.** [35] “Let  $L$  be an ism mapping of  $Q$  into  $H$ . Let  $N_Q$  be the normal cone to  $Q$  at  $z \in Q$ , i.e.  $N_Q z = \{w \in H : \langle z - u, w \rangle \geq 0, \text{ for all } u \in Q\}$ , and define

$$Tz = \begin{cases} Lz + N_Q z, & z \in Q \\ \phi, & z \in H \setminus Q. \end{cases}$$

Then,  $T$  is maximal monotone and  $0 \in Tz$  if and only if  $z \in VI(Q, L)$ .”

**Lemma 2.2.** [36, 38] “Assume that  $H$  is a real Hilbert space. Let mapping  $g : H \rightarrow H$  be Lipschitz continuous monotone and mapping  $A : H \rightarrow 2^H$  be maximal monotone. Then the mapping  $K = g + A$  is maximal monotone.”

**Lemma 2.3.** [32, 37] “Let  $H$  be a real Hilbert space. Then for all  $\zeta, \vartheta \in H$  and  $\gamma \in (0, 1)$ , we have

- (i)  $2\langle \zeta, \vartheta \rangle = \|\zeta\|^2 + \|\vartheta\|^2 - \|\zeta - \vartheta\|^2 = \|\zeta + \vartheta\|^2 - \|\zeta\|^2 - \|\vartheta\|^2.$
- (ii)  $\|\gamma\zeta + (1 - \gamma)\vartheta\|^2 = \gamma\|\zeta\|^2 + (1 - \gamma)\|\vartheta\|^2 - \gamma(1 - \gamma)\|\zeta - \vartheta\|^2.$
- (iii)  $\|\zeta + \vartheta\|^2 \leq \|\zeta\|^2 + 2\langle \vartheta, \zeta + \vartheta \rangle.$

**Lemma 2.4.** [38] “Assume  $H$  is a real Hilbert space,  $B : H \rightarrow H$  is a mapping and  $A : H \rightarrow 2^H$  is maximal monotone mapping. Define the fixed point set of the mapping  $U$  as  $Fix(U) = \{\zeta : \zeta = U\zeta\}$  and  $U_\mu = (I + \mu A)^{-1}(I - \mu B)$ ,  $\mu > 0$ . Then,  $Fix(U_\mu) = (A + B)^{-1}(0)$ , for all  $\mu > 0$ .”

**Lemma 2.5.** [38] “Assume that the sequences  $\{\varphi_n\}$  and  $\{\vartheta_n\}$  are created by the following:

$$\begin{cases} \varphi_n = \zeta_n + \alpha_n(\zeta_n - \zeta_{n-1}); \\ \vartheta_n = (I + \tau A)^{-1}(I - \tau B)\varphi_n. \end{cases}$$

If  $\lim_{n \rightarrow \infty} \|\varphi_n - \vartheta_n\| = 0$  and  $\{\varphi_{n_k}\}$  converges weakly to some  $\iota \in H$ , then  $\iota \in (A + B)^{-1}(0)$ .”

**Lemma 2.6.** [39] “Let  $H$  be a real Hilbert space. Let  $\{\zeta_i, i = 1, 2, \dots, m\} \subset H$ . For  $\{\beta_i\} \subset (0, 1)$ ,  $i = 1, 2, \dots, m$  such that  $\sum_{i=1}^m \beta_i = 1$ , the following identity holds:

$$\left\| \sum_{i=1}^m \beta_i \zeta_i \right\|^2 = \sum_{i=1}^m \beta_i \|\zeta_i\|^2 - \sum_{j,i=1, j \neq i} \beta_i \beta_j \|\zeta_i - \zeta_j\|^2.”$$

**Lemma 2.7.** [25, 40] “Let  $\{\zeta_n\}$  be a sequence of nonnegative real numbers, such that there exists a subsequence  $\{\zeta_{n_j}\}$  of  $\{\zeta_n\}$  such that  $\zeta_{n_j} < \zeta_{n_{j+1}}$  for all  $j \in \mathbb{N}$ . Then, there exists a nondecreasing sequence  $\{n_k\}$  of  $\mathbb{N}$  such that  $\lim_{n \rightarrow \infty} n_k = \infty$  and the following properties are satisfied by all (sufficiently large) number  $k \in \mathbb{N} : \zeta_{n_k} \leq \zeta_{n_{k+1}}$  and  $\zeta_k \leq \zeta_{n_{k+1}}$ . In fact,  $n_k$  is the largest number  $n$  in the set  $\{1, 2, \dots, k\}$ , such that  $\zeta_n < \zeta_{n+1}$ .”

### 3 Main Result

In this section, we introduce a method for finding common solution of variational inequality, finite family of monotone inclusion and fixed point problems of demicontractive mappings in a real Hilbert space. Suppose  $Q$  is nonempty, convex and closed subset of a real Hilbert space  $H$  and

$S_i$  ( $i = 1, 2, \dots, m$ ) is finite family of demicontractive mappings having constant  $k_i \in (0, 1)$  such that  $I - S_i$  are demiclosed at origin. Assume that  $L : Q \rightarrow H$  is  $\sigma$ -inverse strongly monotone mapping,  $A_j : H \rightarrow 2^H$  are maximal monotone mappings,  $B_j : H \rightarrow H$  are  $\xi$ -cocoercive mappings,  $g$  is contractive mapping on  $H$  having constant  $\theta \in (0, 1)$  and constants  $\tau_j$  such that  $0 < \tau_j \leq 2\xi$  for all  $j = 1, 2, \dots, N$ .

**Algorithm 3.1.** Consider  $\beta_{n,i}, \gamma_n \in (0, 1)$  such that  $\sum_{i=1}^m \beta_{n,i} = 1, \beta_{n,0} \in (k, 1)$ ,  $\limsup_{n \rightarrow \infty} \beta_{n,i}(\beta_{n,0} - k) \geq 0, \lim_{n \rightarrow \infty} \gamma_n = 0, \sum_{n=1}^{\infty} \gamma_n = \infty$ , where  $k = \sup_{i \geq 1} \{k_i\} < 1$ . Select  $\eta_n \in (\epsilon, 2\sigma - \epsilon)$ , where  $\epsilon > 0$  and  $\alpha_n \subseteq [0, \infty)$  such that  $\lim_{n \rightarrow \infty} \frac{\alpha_n}{\gamma_n} = 0$ . For  $\zeta_1 \in H$ , calculate  $\{\zeta_{n+1}\}$  using the sequences  $\{\epsilon_n\}, \{\varphi_n\}, \{v_n\}, \{\psi_n\}, \{\kappa_n\}$  as follows:

$$\epsilon_n = \begin{cases} \min \left\{ \frac{\alpha_n}{\|\zeta_n - \zeta_{n-1}\|}, \epsilon \right\}, & \text{if } \zeta_n \neq \zeta_{n-1}, \\ \epsilon, & \text{otherwise} \end{cases}$$

$$\begin{cases} \varphi_n = \zeta_n + \epsilon_n(\zeta_n - \zeta_{n-1}); \\ v_n = \gamma_n g(\varphi_n) + (1 - \gamma_n)\varphi_n; \\ \psi_n = \beta_{n,0}v_n + \sum_{i=1}^m \beta_{n,i}S_i v_n; \\ \kappa_n = P_Q(\psi_n - \eta_n L\psi_n); \\ \zeta_{n+1} = J_{\tau_N}^{A_N}(I - \tau_N B_N)J_{\tau_{N-1}}^{A_{N-1}}(I - \tau_{N-1}B_{N-1})J_{\tau_{N-2}}^{A_{N-2}}(I - \tau_{N-2}B_{N-2}) \dots J_{\tau_1}^{A_1}(I - \tau_1 B_1)\kappa_n. \end{cases} \quad (3.1)$$

Where  $J_{\tau_j}^{A_j}$  ( $j = 1, 2, \dots, N$ ) represents the resolvent mapping of the mapping  $A_j$  and  $I$  represents the identity operator on  $H$ .

**Remark 1.** As  $\epsilon_n \|\zeta_n - \zeta_{n-1}\| \leq \alpha_n$  for all  $n$  and from our assumption  $\lim_{n \rightarrow \infty} \frac{\alpha_n}{\gamma_n} = 0$ , we have

$$\lim_{n \rightarrow \infty} \frac{\epsilon_n}{\gamma_n} \|\zeta_n - \zeta_{n-1}\| \leq \lim_{n \rightarrow \infty} \frac{\alpha_n}{\gamma_n} = 0.$$

Hence,

$$\lim_{n \rightarrow \infty} \frac{\epsilon_n}{\gamma_n} \|\zeta_n - \zeta_{n-1}\| = 0. \quad (3.2)$$

**Theorem 3.2.** Suppose that the solution set  $\Omega = \left\{ \bigcap_{i=1}^m \text{Fix}(S_i) \right\} \cap VI(L, Q) \cap \left\{ \bigcap_{j=1}^N (A_j + B_j)^{-1}(0) \right\}$  is nonempty. Then the sequence  $\{\zeta_n\}$  generated by Algorithm (3.1) converges strongly to an element in  $\Omega$ .

*Proof.* Firstly we will prove  $\{\zeta_n\}$  is bounded. Suppose  $\iota \in \Omega$  and

$\phi^N = J_{\tau_N}^{A_N}(I - \tau_N B_N)J_{\tau_{N-1}}^{A_{N-1}}(I - \tau_{N-1}B_{N-1})J_{\tau_{N-2}}^{A_{N-2}}(I - \tau_{N-2}B_{N-2}) \dots J_{\tau_1}^{A_1}(I - \tau_1 B_1)$ , where  $\phi^0 = I$ . As the resolvent mapping is nonexpansive and using equation (3.1), we have

$$\|\zeta_{n+1} - \iota\|^2 = \|J_{\tau_N}^{A_N}(I - \tau_N B_N)\phi^{N-1}\kappa_n - \iota\|^2 \leq \|\phi^{N-1}\kappa_n - \iota\|^2 \leq \|\phi^{N-2}\kappa_n - \iota\|^2 \leq \|\kappa_n - \iota\|^2. \quad (3.3)$$

Since  $\iota \in VI(L, Q)$ . Also, from equation (3.1) and the fact that projection mapping is nonexpansive, we obtain

$$\|\kappa_n - \iota\|^2 = \|P_Q(\psi_n - \eta_n L\psi_n) - \iota\|^2$$

$$\begin{aligned}
&\leq \|(I - \eta_n L)(\psi_n) - (I - \eta_n L)\iota\|^2 \\
&= \|(\psi_n - \iota) - \eta_n(L\psi_n - L\iota)\|^2 \\
&= \|\psi_n - \iota\|^2 + \eta_n^2 \|L\psi_n - L\iota\|^2 - 2\eta_n \langle \psi_n - \iota, L\psi_n - L\iota \rangle \\
&\leq \|\psi_n - \iota\|^2 + \eta_n^2 \|L\psi_n - L\iota\|^2 - 2\eta_n \sigma \|L\psi_n - L\iota\|^2 \\
&= \|\psi_n - \iota\|^2 - \eta_n(2\sigma - \eta_n) \|L\psi_n - L\iota\|^2.
\end{aligned} \tag{3.4}$$

Since  $\eta_n \in (\epsilon, 2\sigma - \epsilon)$ , so we have

$$\|\kappa_n - \iota\|^2 \leq \|\psi_n - \iota\|^2. \tag{3.5}$$

Using Lemmas (2.3), (2.6) and equation (3.1), we have

$$\begin{aligned}
\|\psi_n - \iota\|^2 &\leq \left\| \beta_{n,0}v_n + \sum_{i=1}^m \beta_{n,i}S_i v_n - \iota \right\|^2 \\
&= \left\| \beta_{n,0}(v_n - \iota) + \sum_{i=1}^m \beta_{n,i}(S_i v_n - \iota) \right\|^2 \\
&= \beta_{n,0}\|v_n - \iota\|^2 + \sum_{i=1}^m \beta_{n,i}\|S_i v_n - \iota\|^2 - \sum_{i=1}^m \beta_{n,0}\beta_{n,i}\|v_n - S_i v_n\|^2 \\
&\leq \beta_{n,0}\|v_n - \iota\|^2 + \sum_{i=1}^m \beta_{n,i} [\|v_n - \iota\|^2 + k\|v_n - S_i v_n\|^2] - \sum_{i=1}^m \beta_{n,0}\beta_{n,i}\|v_n - S_i v_n\|^2 \\
&= \|v_n - \iota\|^2 + (k - \beta_{n,0}) \sum_{i=1}^m \beta_{n,i}\|v_n - S_i v_n\|^2.
\end{aligned} \tag{3.6}$$

Since  $\beta_{n,0} \in (k, 1)$ . From equation (3.6), we have

$$\|\psi_n - \iota\|^2 \leq \|v_n - \iota\|^2. \tag{3.7}$$

Again from equation (3.1), we have

$$\begin{aligned}
\|\varphi_n - \iota\| &= \|\zeta_n + \epsilon_n(\zeta_n - \zeta_{n-1}) - \iota\| \\
&\leq \|\zeta_n - \iota\| + \gamma_n \frac{\epsilon_n}{\gamma_n} \|\zeta_n - \zeta_{n-1}\|.
\end{aligned} \tag{3.8}$$

By Remark (1), we have  $\lim_{n \rightarrow \infty} \frac{\epsilon_n}{\gamma_n} \|\zeta_n - \zeta_{n-1}\| = 0$ , therefore there exist a constant  $M_1 > 0$  such that  $\frac{\epsilon_n}{\gamma_n} \|\zeta_n - \zeta_{n-1}\| \leq M_1$ . Therefore, equation (3.8) implies

$$\|\varphi_n - \iota\| \leq \|\zeta_n - \iota\| + \gamma_n M_1. \tag{3.9}$$

Using equations (3.3), (3.5), (3.7) and (3.9) we deduce that

$$\begin{aligned}
\|\zeta_{n+1} - \iota\| &\leq \|v_n - \iota\| \\
&= \|\gamma_n g(\varphi_n) + (1 - \gamma_n)\varphi_n - \iota\| \\
&= \|\gamma_n(g(\varphi_n) - \iota) + (1 - \gamma_n)(\varphi_n - \iota)\| \\
&= \|\gamma_n(g(\varphi_n) - g(\iota)) + \gamma_n(g(\iota) - \iota) + (1 - \gamma_n)(\varphi_n - \iota)\| \\
&\leq \gamma_n \|g(\varphi_n) - g(\iota)\| + \gamma_n \|g(\iota) - \iota\| + (1 - \gamma_n) \|\varphi_n - \iota\|
\end{aligned} \tag{3.10}$$

$$\begin{aligned}
&\leq \gamma_n \theta \|\varphi_n - \iota\| + (1 - \gamma_n) \|\varphi_n - \iota\| + \gamma_n \|g(\iota) - \iota\| \\
&= [1 - \gamma_n(1 - \theta)] \|\varphi_n - \iota\| + \gamma_n \|g(\iota) - \iota\| \\
&\leq [1 - \gamma_n(1 - \theta)] [\|\zeta_n - \iota\| + \gamma_n M_1] + \gamma_n(1 - \theta) \left( \frac{\|g(\iota) - \iota\|}{1 - \theta} \right) \\
&\leq [1 - \gamma_n(1 - \theta)] [\|\zeta_n - \iota\| + M_1] + \gamma_n(1 - \theta) \left( \frac{\|g(\iota) - \iota\|}{1 - \theta} \right).
\end{aligned}$$

Continuing like this,

$$\|\zeta_{n+1} - \iota\| \leq \max \left\{ \|\zeta_0 - \iota\| + M_1, \frac{\|g(\iota) - \iota\|}{1 - \theta} \right\}.$$

Thus  $\{\zeta_n\}$  is bounded and hence  $\{\psi_n\}$ ,  $\{\varphi_n\}$ ,  $\{v_n\}$  and  $\{\kappa_n\}$  are also bounded. Now, using equation (3.1) and Lemma (2.3), we obtain

$$\begin{aligned}
\|v_n - \iota\|^2 &= \|\gamma_n g(\varphi_n) + (1 - \gamma_n) \varphi_n - \iota\|^2 \\
&= \|\gamma_n (g(\varphi_n) - \iota) + (1 - \gamma_n) (\varphi_n - \iota)\|^2 \\
&\leq (1 - \gamma_n) \|\varphi_n - \iota\|^2 + 2\gamma_n \langle g(\varphi_n) - \iota, \varphi_n - \iota + \gamma_n (g(\varphi_n) - \varphi_n) \rangle.
\end{aligned} \tag{3.11}$$

From equations (3.3), (3.5), (3.6) and (3.11), we obtain

$$\begin{aligned}
\|\zeta_{n+1} - \iota\|^2 &\leq \|\psi_n - \iota\|^2 \\
&\leq \|v_n - \iota\|^2 + (k - \beta_{n,0}) \sum_{i=1}^m \beta_{n,i} \|v_n - S_i v_n\|^2 \\
&\leq (1 - \gamma_n) \|\varphi_n - \iota\|^2 + 2\gamma_n \langle g(\varphi_n) - \iota, \varphi_n - \iota + \gamma_n (g(\varphi_n) - \varphi_n) \rangle \\
&\quad + (k - \beta_{n,0}) \sum_{i=1}^m \beta_{n,i} \|v_n - S_i v_n\|^2.
\end{aligned} \tag{3.12}$$

Since,  $\beta_{n,0} \in (k, 1)$ , so equation (3.12) gives

$$\|\zeta_{n+1} - \iota\|^2 \leq (1 - \gamma_n) \|\varphi_n - \iota\|^2 + 2\gamma_n \langle g(\varphi_n) - \iota, \varphi_n - \iota + \gamma_n (g(\varphi_n) - \varphi_n) \rangle. \tag{3.13}$$

Now, from equation (3.1), we have

$$\begin{aligned}
\|\varphi_n - \iota\|^2 &= \|\zeta_n + \epsilon_n (\zeta_n - \zeta_{n-1}) - \iota\|^2 \\
&= \|\zeta_n - \iota\|^2 + \epsilon_n^2 \|\zeta_n - \zeta_{n-1}\|^2 + 2\epsilon_n \langle \zeta_n - \iota, \zeta_n - \zeta_{n-1} \rangle \\
&\leq \|\zeta_n - \iota\|^2 + \epsilon_n^2 \|\zeta_n - \zeta_{n-1}\|^2 + 2\epsilon_n \|\zeta_n - \iota\| \|\zeta_n - \zeta_{n-1}\|.
\end{aligned} \tag{3.14}$$

Now, we prove  $\zeta_n \rightarrow \iota$ . For this, we study two cases.

**Case 1:** Suppose that there exists a number  $N \in \mathbb{N}$  such that  $\|\zeta_{n+1} - \iota\|^2 \leq \|\zeta_n - \iota\|^2$  for any  $n \geq N$ . As sequence  $\{\|\zeta_n - \iota\|\}^2$  is bounded and monotonic, this gives  $\{\|\zeta_n - \iota\|\}^2$  is convergent. From equations (3.12) and (3.14), we have

$$\begin{aligned}
(\beta_{n,0} - k) \sum_{i=1}^m \beta_{n,i} \|v_n - S_i v_n\|^2 &\leq (1 - \gamma_n) \{ \|\zeta_n - \iota\|^2 + \epsilon_n^2 \|\zeta_n - \zeta_{n-1}\|^2 + 2\epsilon_n \|\zeta_n - \iota\| \|\zeta_n - \zeta_{n-1}\| \} \\
&\quad + 2\gamma_n \langle g(\varphi_n) - \iota, \varphi_n - \iota + \gamma_n (g(\varphi_n) - \varphi_n) \rangle - \|\zeta_{n+1} - \iota\|^2
\end{aligned}$$

$$\begin{aligned} &\leq \|\zeta_n - \iota\|^2 + \epsilon_n^2 \|\zeta_n - \zeta_{n-1}\|^2 + 2\epsilon_n \|\zeta_n - \iota\| \|\zeta_n - \zeta_{n-1}\| \\ &+ 2\gamma_n \langle g(\varphi_n) - \iota, \varphi_n - \iota + \gamma_n(g(\varphi_n) - \varphi_n) \rangle - \|\zeta_{n+1} - \iota\|^2, \end{aligned}$$

which implies

$$\begin{aligned} (\beta_{n,0} - k) \sum_{i=1}^m \beta_{n,i} \|v_n - S_i v_n\|^2 &\leq \|\zeta_n - \iota\|^2 + \frac{\epsilon_n^2}{\gamma_n^2} \gamma_n^2 \|\zeta_n - \zeta_{n-1}\|^2 + \frac{2\epsilon_n}{\gamma_n} \gamma_n \|\zeta_n - \iota\| \|\zeta_n - \zeta_{n-1}\| \\ &+ 2\gamma_n \langle g(\varphi_n) - \iota, \varphi_n - \iota + \gamma_n(g(\varphi_n) - \varphi_n) \rangle - \|\zeta_{n+1} - \iota\|^2. \end{aligned} \quad (3.15)$$

Taking limit  $n \rightarrow \infty$  in equation (3.15), by using Remark (1) and  $\lim_{n \rightarrow \infty} \gamma_n = 0$ , we obtain

$$\lim_{n \rightarrow \infty} \|v_n - S_i v_n\| = 0. \quad (3.16)$$

Using equation (3.1), we have

$$\begin{aligned} \|v_n - \varphi_n\| &= \|\gamma_n g(\varphi_n) + (1 - \gamma_n)\varphi_n - \varphi_n\| \\ &= \gamma_n \|g(\varphi_n) - \varphi_n\|. \end{aligned} \quad (3.17)$$

Taking limit  $n \rightarrow \infty$  in equation (3.17) and using  $\lim_{n \rightarrow \infty} \gamma_n = 0$ , we obtain

$$\lim_{n \rightarrow \infty} \|v_n - \varphi_n\| = 0. \quad (3.18)$$

From equation (3.1), we have

$$\|\psi_n - v_n\| \leq \sum_{i=1}^m \beta_{n,i} \|S_i v_n - v_n\|. \quad (3.19)$$

Taking limit  $n \rightarrow \infty$  in above equation and using equation (3.16), we get

$$\lim_{n \rightarrow \infty} \|\psi_n - v_n\| = 0. \quad (3.20)$$

Using triangle inequality, equations (3.18) and (3.20), we have

$$\lim_{n \rightarrow \infty} \|\psi_n - \varphi_n\| = 0. \quad (3.21)$$

From equations (3.3), (3.4) and (3.7), we have

$$\begin{aligned} \|\zeta_{n+1} - \iota\|^2 &\leq \|\kappa_n - \iota\|^2 \\ &\leq \|\psi_n - \iota\|^2 - \eta_n(2\sigma - \eta_n) \|L\psi_n - L\iota\|^2 \\ &\leq \|v_n - \iota\|^2 - \eta_n(2\sigma - \eta_n) \|L\psi_n - L\iota\|^2, \end{aligned} \quad (3.22)$$

which implies

$$\eta_n(2\sigma - \eta_n) \|L\psi_n - L\iota\|^2 \leq \|v_n - \iota\|^2 - \|\zeta_{n+1} - \iota\|^2. \quad (3.23)$$

Using equations (3.11) and (3.23), we deduce

$$\eta_n(2\sigma - \eta_n) \|L\psi_n - L\iota\|^2 \leq (1 - \gamma_n) \|\varphi_n - \iota\|^2 + 2\gamma_n \langle g(\varphi_n) - \iota, \varphi_n - \iota + \gamma_n(g(\varphi_n) - \varphi_n) \rangle - \|\zeta_{n+1} - \iota\|^2. \quad (3.24)$$

Consequently, from equation (3.14) and (3.24), we have

$$\begin{aligned} \eta_n(2\sigma - \eta_n) \|L\psi_n - L\iota\|^2 &\leq \|\zeta_n - \iota\|^2 - \|\zeta_{n+1} - \iota\|^2 + 2\gamma_n \langle g(\varphi_n) - \iota, \varphi_n - \iota + \gamma_n(g(\varphi_n) - \varphi_n) \rangle \\ &+ \epsilon_n^2 \|\zeta_n - \zeta_{n-1}\|^2 + 2\epsilon_n \|\zeta_n - \iota\| \|\zeta_n - \zeta_{n-1}\|. \end{aligned} \quad (3.25)$$

Taking limit  $n \rightarrow \infty$  in equation (3.25) and using Remark (1) and  $\lim_{n \rightarrow \infty} \gamma_n = 0$ , we have

$$\lim_{n \rightarrow \infty} \|L\psi_n - L\iota\| = 0. \quad (3.26)$$



Using equation (3.1) and Lemma (2.3), we obtain

$$\begin{aligned} \|\kappa_n - \iota\|^2 &= \|P_C(\psi_n - \eta_n L\psi_n) - \iota\|^2 \\ &\leq \langle \kappa_n - \iota, (I - \eta_n L)\psi_n - (I - \eta_n L)\iota \rangle \\ &= \frac{1}{2} \{ \|\kappa_n - \iota\|^2 + \|(I - \eta_n L)\psi_n - (I - \eta_n L)\iota\|^2 - \|\kappa_n - \psi_n + \eta_n(L\psi_n - L\iota)\|^2 \}. \end{aligned}$$

After rearranging the terms, we have

$$\|\kappa_n - \iota\|^2 \leq \|\psi_n - \iota\|^2 - \|\kappa_n - \psi_n\|^2 - \eta_n^2 \|L\psi_n - L\iota\|^2 - 2\eta_n \|\kappa_n - \psi_n\| \|L\psi_n - L\iota\|. \quad (3.27)$$

Now, from equation (3.13) and (3.14), we have

$$\begin{aligned} \|\psi_n - \iota\|^2 &\leq (1 - \gamma_n) \|\varphi_n - \iota\|^2 + 2\gamma_n \langle g(\varphi_n) - \iota, \varphi_n - \iota + \gamma_n(g\varphi_n - \varphi_n) \rangle \\ &\leq (1 - \gamma_n) \{ \|\zeta_n - \iota\|^2 + \epsilon_n^2 \|\zeta_n - \zeta_{n-1}\|^2 + 2\epsilon_n \|\zeta_n - \iota\| \|\zeta_n - \zeta_{n-1}\| \} \\ &\quad + 2\gamma_n \langle g(\varphi_n) - \iota, \varphi_n - \iota + \gamma_n(g(\varphi_n) - \varphi_n) \rangle. \end{aligned} \quad (3.28)$$

Again, using equations (3.3), (3.27) and (3.28), we deduce

$$\begin{aligned} \|\zeta_{n+1} - \iota\|^2 &\leq \|\kappa_n - \iota\|^2 \\ &\leq (1 - \gamma_n) \{ \|\zeta_n - \iota\|^2 + \epsilon_n^2 \|\zeta_n - \zeta_{n-1}\|^2 + 2\epsilon_n \|\zeta_n - \iota\| \|\zeta_n - \zeta_{n-1}\| \} - \|\kappa_n - \psi_n\|^2 \\ &\quad + 2\eta_n \|\kappa_n - \psi_n\| \|L\psi_n - L\iota\| \\ &\quad + 2\gamma_n \langle g(\varphi_n) - \iota, \varphi_n - \iota + \gamma_n(g(\varphi_n) - \varphi_n) \rangle - \eta_n^2 \|L\psi_n - L\iota\|^2 \\ &\leq \|\zeta_n - \iota\|^2 + \epsilon_n^2 \|\zeta_n - \zeta_{n-1}\|^2 + 2\epsilon_n \|\zeta_n - \iota\| \|\zeta_n - \zeta_{n-1}\| - \|\kappa_n - \psi_n\|^2 \\ &\quad + 2\eta_n \|\psi_n - \iota\| \|L\psi_n - L\iota\| \\ &\quad + 2\gamma_n \langle g(\varphi_n) - \iota, \varphi_n - \iota + \gamma_n(g(\varphi_n) - \varphi_n) \rangle - \eta_n^2 \|L\psi_n - L\iota\|^2, \end{aligned}$$

which implies

$$\begin{aligned} \|\kappa_n - \psi_n\|^2 &\leq \|\zeta_n - \iota\|^2 - \|\zeta_{n+1} - \iota\|^2 + 2\eta_n \|\psi_n - \iota\| \|L\psi_n - L\iota\| - \eta_n^2 \|L\psi_n - L\iota\|^2 \\ &\quad + 2\gamma_n \langle g(\varphi_n) - \iota, \varphi_n - \iota + \gamma_n(g(\varphi_n) - \varphi_n) \rangle \\ &\quad + \epsilon_n^2 \|\zeta_n - \zeta_{n-1}\|^2 + 2\epsilon_n \|\zeta_n - \iota\| \|\zeta_n - \zeta_{n-1}\|. \end{aligned} \quad (3.29)$$

Taking limit  $n \rightarrow \infty$  in equation (3.29) and using equation (3.26), remark (1) and  $\lim_{n \rightarrow \infty} \gamma_n = 0$ , we get

$$\lim_{n \rightarrow \infty} \|\kappa_n - \psi_n\| = 0. \quad (3.30)$$

From equations (3.21), (3.30) and triangle inequality, we have

$$\lim_{n \rightarrow \infty} \|\kappa_n - \varphi_n\| = 0. \quad (3.31)$$

Consider

$$\|\zeta_n - \varphi_n\| = \|\epsilon_n(\zeta_n - \zeta_{n-1})\|. \quad (3.32)$$

By using Remark (1) and taking limit  $n \rightarrow \infty$  in equation (3.32), we get

$$\lim_{n \rightarrow \infty} \|\zeta_n - \varphi_n\| = 0. \quad (3.33)$$

By using equations (3.31), (3.33) and triangle inequality, we deduce

$$\lim_{n \rightarrow \infty} \|\kappa_n - \zeta_n\| = 0. \quad (3.34)$$

From equations (3.30) and (3.34), we have

$$\lim_{n \rightarrow \infty} \|\psi_n - \zeta_n\| = 0. \quad (3.35)$$

Again using equations (3.20), (3.30) and (3.34), we get

$$\lim_{n \rightarrow \infty} \|\zeta_n - v_n\| = 0. \quad (3.36)$$

Now, consider

$$\begin{aligned} \|\zeta_{n+1} - \iota\|^2 &= \|J_{\tau_N}^{A_N}(I - \tau_N B_N)J_{\tau_N}^{A_{N-1}}(I - \tau_{N-1}B_{N-1})J_{\tau_{N-2}}^{A_{N-2}}(I - \tau_{N-2}B_{N-2})\dots J_{\tau_1}^{A_1}(I - \tau_1 B_1)\kappa_n - \iota\|^2 \\ &= \|J_{\tau_N}^{A_N}(I - \tau_N B_N)\phi^{N-1}\kappa_n - \iota\|^2. \end{aligned}$$

As  $J_{\tau_N}^{A_N}(I - \tau_N B_N)$  is firmly nonexpansive, we have

$$\|\zeta_{n+1} - \iota\|^2 \leq \langle \zeta_{n+1} - \iota, \phi^{N-1}\kappa_n - \iota \rangle. \quad (3.37)$$

Using Lemma (2.3) and equation (3.37), we obtain

$$\|\zeta_{n+1} - \iota\|^2 \leq \frac{1}{2} \left[ \|\zeta_{n+1} - \iota\|^2 + \|\phi^{N-1}\kappa_n - \iota\|^2 - \|\zeta_{n+1} - \phi^{N-1}\kappa_n\|^2 \right]. \quad (3.38)$$

From equation (3.38), we have

$$\|\zeta_{n+1} - \iota\|^2 \leq \|\phi^{N-1}\kappa_n - \iota\|^2 - \|\zeta_{n+1} - \phi^{N-1}\kappa_n\|^2,$$

which implies

$$\|\zeta_{n+1} - \phi^{N-1}\kappa_n\|^2 \leq \|\phi^{N-1}\kappa_n - \iota\|^2 - \|\zeta_{n+1} - \iota\|^2. \quad (3.39)$$

Taking limit  $n \rightarrow \infty$  in equation (3.39) and using equations (3.3) and (3.34), we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \|\zeta_{n+1} - \phi^{N-1}\kappa_n\|^2 &\leq \lim_{n \rightarrow \infty} \left( \|\phi^{N-1}\kappa_n - \iota\|^2 - \|\zeta_{n+1} - \iota\|^2 \right) \\ &\leq \lim_{n \rightarrow \infty} \left( \|\kappa_n - \iota\|^2 - \|\zeta_{n+1} - \iota\|^2 \right) \\ &= \lim_{n \rightarrow \infty} \left( \|\kappa_n - \zeta_n\|^2 + 2\|\kappa_n - \zeta_n\|\|\zeta_n - \iota\| \right. \\ &\quad \left. + \|\zeta_n - \iota\|^2 - \|\zeta_{n+1} - \iota\|^2 \right) \\ &\leq \lim_{n \rightarrow \infty} \left( \|\kappa_n - \zeta_n\|^2 + 2\|\kappa_n - \zeta_n\|\|\zeta_n - \iota\| \right) \\ &\quad + \lim_{n \rightarrow \infty} \left( \|\zeta_n - \iota\|^2 - \|\zeta_{n+1} - \iota\|^2 \right) \\ &\leq 0. \end{aligned}$$

Therefore,

$$\lim_{n \rightarrow \infty} \|\zeta_{n+1} - \phi^{N-1}\kappa_n\| = 0. \quad (3.40)$$

Using the similar argument as in (3.39), we get

$$\|\phi^{N-1}\kappa_n - \phi^{N-2}\kappa_n\|^2 \leq \left( \|\phi^{N-2}\kappa_n - \iota\|^2 - \|\phi^{N-1}\kappa_n - \iota\|^2 \right). \quad (3.41)$$

Taking limit  $n \rightarrow \infty$  in equation (3.41) and using equations (3.3), (3.34) and (3.40), we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \|\phi^{N-1}\kappa_n - \phi^{N-2}\kappa_n\|^2 &\leq \lim_{n \rightarrow \infty} \left( \|\phi^{N-2}\kappa_n - \iota\|^2 - \|\phi^{N-1}\kappa_n - \iota\|^2 \right) \\ &\leq \lim_{n \rightarrow \infty} \left( \|\kappa_n - \iota\|^2 - \|\zeta_{n+1} - \iota\|^2 \right) \\ &= \lim_{n \rightarrow \infty} \left( \|\kappa_n - \zeta_n\|^2 + 2\|\kappa_n - \zeta_n\|\|\zeta_n - \iota\| \right. \\ &\quad \left. + \|\zeta_n - \iota\|^2 - \|\zeta_{n+1} - \iota\|^2 \right) \\ &\leq \lim_{n \rightarrow \infty} \left( \|\kappa_n - \zeta_n\|^2 + 2\|\kappa_n - \zeta_n\|\|\zeta_n - \iota\| \right) \end{aligned}$$

$$\begin{aligned} & + \lim_{n \rightarrow \infty} (\|\zeta_n - \iota\|^2 - \|\zeta_{n+1} - \iota\|^2) \\ & \leq 0. \end{aligned} \tag{3.42}$$

Hence,  $\lim_{n \rightarrow \infty} \|\phi^{N-1}\kappa_n - \phi^{N-2}\kappa_n\| = 0$ .

Continuing like this,

$$\lim_{n \rightarrow \infty} \|\phi^j \kappa_n - \phi^{j-1} \kappa_n\| = 0 \quad \text{for all } j = 1, 2, \dots, N. \tag{3.43}$$

Also, using triangle inequality, we obtain

$$\|\zeta_{n+1} - \kappa_n\| \leq \|\phi^N \kappa_n - \phi^{N-1} \kappa_n\| + \|\phi^{N-1} \kappa_n - \phi^{N-2} \kappa_n\| + \dots + \|\phi^1 \kappa_n - \kappa_n\|. \tag{3.44}$$

Taking limit  $n \rightarrow \infty$  in equation (3.44) and using equation (3.43), we have

$$\lim_{n \rightarrow \infty} \|\zeta_{n+1} - \kappa_n\| = 0. \tag{3.45}$$

From equations (3.30), (3.35) and (3.45), we obtain

$$\lim_{n \rightarrow \infty} \|\zeta_{n+1} - \zeta_n\| = 0. \tag{3.46}$$

As the sequence  $\{\zeta_n\}$  is bounded, so there exists a subsequence  $\{\zeta_{n_k}\}$  of  $\{\zeta_n\}$  such that  $\{\zeta_{n_k}\}$  converges to  $q$ . Using equations (3.35) and (3.36), there exists subsequences  $\{v_{n_k}\}$  of  $\{v_n\}$  and  $\{\psi_{n_k}\}$  of  $\{\psi_n\}$  that converges weakly to  $q$ . Hence, by demiclosedness of  $S_i$  at zero and using equation (3.16), we have  $q \in \text{Fix}(S_i)$  for each  $i = 1, 2, \dots, m$ . We next show that  $q \in VI(L, Q)$ . Let the mapping  $M'$  be defined by

$$M'(z^*) = \begin{cases} L(z^*) + N_Q(z^*) & \text{if } z^* \in Q \\ \phi & \text{if } z^* \notin Q \end{cases},$$

where  $N_Q(z^*) = \{t' \in H : \langle z^* - p', t' \rangle \geq 0 \text{ for all } p' \in Q\}$  is the normal cone at  $Q$  and  $M'$  is maximal monotone, which implies  $0 \in M'(z^*)$  if and only if  $z^* \in VI(L, Q)$ . Let us assume that  $(z^*, t') \in G(M')$ , where  $G(M')$  denotes graph of mapping  $M'$ . So,  $t' \in M'(z^*) = L(z^*) + N_Q(z^*)$  which implies  $t' - L(z^*) \in N_Q(z^*)$ .

Hence, we obtain  $\langle z^* - p', t' - L(z^*) \rangle \geq 0$  for all  $p' \in Q$ . Also, from equation (3.1) and using that  $z^* \in Q$ , we get  $\langle \psi_n - \eta_n L\psi_n - \kappa_n, \kappa_n - z^* \rangle \geq 0$ . Hence,  $\langle z^* - \kappa_n, \frac{\kappa_n - \psi_n}{\eta_n} + L\psi_n \rangle \geq 0$ . As  $\langle z^* - p, t' - Lz^* \rangle \geq 0$  for all  $p' \in Q, \kappa_n \in Q$  and monotonicity of  $L$ , we have

$$\begin{aligned} \langle z^* - \kappa_n, t' \rangle & \geq \langle z^* - \kappa_n, Lz^* \rangle \\ & \geq \langle z^* - \kappa_n, Lz^* \rangle - \langle z^* - \kappa_n, \frac{\kappa_n - \psi_n}{\eta_n} + L\psi_n \rangle \\ & = \langle z^* - \kappa_n, Lz^* - L\kappa_n \rangle - \langle z^* - \kappa_n, \frac{\kappa_n - \psi_n}{\eta_n} \rangle + \langle z^* - \kappa_n, L\kappa_n - L\psi_n \rangle \\ & \geq \langle z^* - \kappa_n, L\kappa_n - L\psi_n \rangle - \langle z^* - \kappa_n, \frac{\kappa_n - \psi_n}{\eta_n} \rangle. \end{aligned}$$

As  $L$  is continuous, taking limit  $n \rightarrow \infty$ , we obtain  $\langle z^* - q, t' \rangle \geq 0$ . Since,  $M'$  is maximal monotone, so we obtain  $q \in M'^{-1}(0)$  and therefore  $0 \in M'(q)$  and hence  $q \in VI(L, Q)$ . Now, we prove  $q \in (A_j + B_j)^{-1}(0)$ . Let us denote  $T_j = (I + \tau_j A_j)^{-1}(I - \tau_j B_j)$  for all  $j = 1, 2, \dots, N$ . Putting  $j = 1$  in equation (3.43), we obtain

$$\lim_{n \rightarrow \infty} \|\phi^1 \kappa_n - \kappa_n\| = 0, \tag{3.47}$$

which implies

$$\lim_{n \rightarrow \infty} \|T_1 \kappa_n - \kappa_n\| = 0. \tag{3.48}$$

As  $T_1$  is nonexpansive, we have

$$\begin{aligned}
\|T_1\varphi_n - \varphi_n\| &= \|T_1\varphi_n - T_1\kappa_n + T_1\kappa_n - \varphi_n\| \\
&\leq \|T_1\varphi_n - T_1\kappa_n\| + \|T_1\kappa_n - \varphi_n\| \\
&\leq \|\varphi_n - \kappa_n\| + \|T_1\kappa_n - \varphi_n\| \\
&\leq \|\varphi_n - \kappa_n\| + \|T_1\kappa_n - \kappa_n + \kappa_n - \varphi_n\| \\
&\leq \|\varphi_n - \kappa_n\| + \|T_1\kappa_n - \kappa_n\| + \|\kappa_n - \varphi_n\|.
\end{aligned} \tag{3.49}$$

Taking limit  $n \rightarrow \infty$  in equation (3.49), we get

$$\lim_{n \rightarrow \infty} \|T_1\varphi_n - \varphi_n\| = 0. \tag{3.50}$$

From equation (3.50) and Lemmas (2.4) and (2.5), we deduce that

$$q \in (A_1 + B_1)^{-1}(0) = \text{Fix}(T_1).$$

Putting  $j = 2$  in equation (3.43), we get

$$\lim_{n \rightarrow \infty} \|\phi^2\kappa_n - \phi^1\kappa_n\| = 0,$$

which implies

$$\lim_{n \rightarrow \infty} \|T_2T_1\kappa_n - T_1\kappa_n\| = 0. \tag{3.51}$$

Consider

$$\begin{aligned}
\|T_2\kappa_n - \kappa_n\| &= \|T_2\kappa_n - T_2T_1\kappa_n + T_2T_1\kappa_n - \kappa_n\| \\
&\leq \|T_2\kappa_n - T_2T_1\kappa_n\| + \|T_2T_1\kappa_n - \kappa_n\| \\
&\leq \|\kappa_n - T_1\kappa_n\| + \|T_2T_1\kappa_n - \kappa_n\| \\
&= \|\kappa_n - T_1\kappa_n\| + \|T_2T_1\kappa_n - T_1\kappa_n + T_1\kappa_n - \kappa_n\| \\
&\leq \|\kappa_n - T_1\kappa_n\| + \|T_2T_1\kappa_n - T_1\kappa_n\| + \|T_1\kappa_n - \kappa_n\|.
\end{aligned} \tag{3.52}$$

Taking limit  $n \rightarrow \infty$  in equation (3.52) and using equations (3.48) and (3.51), we have

$$\lim_{n \rightarrow \infty} \|T_2\kappa_n - \kappa_n\| = 0. \tag{3.53}$$

Similarly, as equation (3.50), we obtain

$$\lim_{n \rightarrow \infty} \|T_2\varphi_n - \varphi_n\| = 0. \tag{3.54}$$

From equation (3.54), Lemmas (2.4) and (2.5), we obtain

$$q \in \text{Fix}(T_2) = (A_2 + B_2)^{-1}(0).$$

Continuing like this, we have  $q \in \text{Fix}(T_j)$  for all  $j = 1, 2, \dots, N$ . Hence,

$$q \in \bigcap_{j=1}^N \text{Fix}(T_j) = \bigcap_{j=1}^N (A_j + B_j)^{-1}(0).$$

From equations (3.13) and (3.14), we obtain

$$\begin{aligned}
\|\zeta_{n+1} - \iota\|^2 &\leq (1 - \gamma_n)\{\|\zeta_n - \iota\|^2 + \epsilon_n^2\|\zeta_n - \zeta_{n-1}\|^2 + 2\epsilon_n\|\zeta_n - \iota\|\|\zeta_n - \zeta_{n-1}\|\} \\
&\quad + 2\gamma_n\langle g(\varphi_n) - \iota, \varphi_n - \iota + \gamma_n(g(\varphi_n) - \varphi_n) \rangle.
\end{aligned} \tag{3.55}$$

Hence

$$\begin{aligned} \|\zeta_{n+1} - \iota\|^2 &\leq (1 - \gamma_n)\|\zeta_n - \iota\|^2 + \gamma_n\{2\langle g(\varphi_n) - \iota, \varphi_n - \iota + \gamma_n(g(\varphi_n) - \varphi_n) \rangle + \frac{\epsilon_n^2 \gamma_n}{\gamma_n^2} \|\zeta_n - \zeta_{n-1}\|^2 \\ &\quad + \frac{2\epsilon_n}{\gamma_n} \|\zeta_n - \iota\| \|\zeta_n - \zeta_{n-1}\|\}, \end{aligned} \quad (3.56)$$

which gives

$$r_{n+1} \leq (1 - \gamma_n)r_n + \gamma_n s_n, \quad (3.57)$$

where  $r_n = \|\zeta_n - \iota\|^2$  and

$$s_n = 2\langle g(\varphi_n) - \iota, \varphi_n - \iota + \gamma_n(g(\varphi_n) - \varphi_n) \rangle + \frac{\epsilon_n^2 \gamma_n}{\gamma_n^2} \|\zeta_n - \zeta_{n-1}\|^2 + \frac{2\epsilon_n}{\gamma_n} \|\zeta_n - \iota\| \|\zeta_n - \zeta_{n-1}\|.$$

Now, it remains to prove that  $\limsup_{n \rightarrow \infty} \langle g(\varphi_n) - \iota, \varphi_n - \iota + \gamma_n(g(\varphi_n) - \varphi_n) \rangle \leq 0$ .

Let  $\{\varphi_{n_k}\}$  be a subsequence of  $\{\varphi_n\}$ . Using equation (3.33),  $\{\varphi_{n_k}\}$  weakly converges to  $q$ . Consider

$$\begin{aligned} \limsup_{n \rightarrow \infty} \langle g(\varphi_n) - \iota, \varphi_n - \iota + \gamma_n(g(\varphi_n) - \varphi_n) \rangle &\leq \limsup_{n \rightarrow \infty} \langle g(\varphi_n) - \iota, \varphi_n - \iota \rangle \\ &\quad + \limsup_{n \rightarrow \infty} \langle g(\varphi_n) - \iota, \gamma_n(g(\varphi_n) - \varphi_n) \rangle \\ &\leq \limsup_{n \rightarrow \infty} \langle g(\varphi_n) - \iota, \varphi_n - \iota \rangle \\ &\quad + \limsup_{n \rightarrow \infty} \gamma_n \langle g(\varphi_n) - \iota, g(\varphi_n) - \varphi_n \rangle \\ &\leq \limsup_{n \rightarrow \infty} \langle g(\varphi_n) - \iota, \varphi_n - \iota \rangle \\ &\leq \langle g(q) - \iota, q - \iota \rangle \leq 0. \end{aligned}$$

Hence,

$$\limsup_{n \rightarrow \infty} \langle g(\varphi_n) - \iota, \varphi_n - \iota + \gamma_n(g(\varphi_n) - \varphi_n) \rangle \leq 0. \quad (3.58)$$

Using equation (3.58), Remark (1) and Lemma (2.1),  $\{r_n\}$  converges to zero as  $n \rightarrow \infty$ . Thus  $\zeta_n \rightarrow \iota$ .

**Case 2:** Suppose that there exists a subsequence  $\{r_{n_j}\}$  of  $\{r_n\}$  such that  $r_{n_j} \leq r_{n_j+1}$  for any  $j \in \mathbb{N}$ . From Lemma (2.7), we observe that there exists a nondecreasing sequence  $\{n_k\}$  of  $\mathbb{N}$  such that  $\lim_{k \rightarrow \infty} n_k = \infty$  which satisfy following inequalities for all  $k \in \mathbb{N}$ .

$$r_{n_k} \leq r_{n_k+1} \quad (3.59)$$

and

$$r_k \leq r_{n_k+1}. \quad (3.60)$$

Using equation (3.3), we obtain

$$\begin{aligned} \|v_{n_k} - \rho_{n_k}\| &= \|\gamma_{n_k} g(\rho_{n_k}) + (1 - \gamma_{n_k})\rho_{n_k} - \rho_{n_k}\| \\ &= \gamma_{n_k} \|g(\rho_{n_k}) - \rho_{n_k}\|. \end{aligned} \quad (3.61)$$

Taking limit  $k \rightarrow \infty$  in equation (3.61), we get

$$\lim_{n \rightarrow \infty} \|v_{n_k} - \rho_{n_k}\| = 0. \quad (3.62)$$

From equation (3.19), we have

$$\|\psi_{n_k} - v_{n_k}\| \leq \sum_{i=1}^m \beta_{n_k, i} \|S_i v_{n_k} - v_{n_k}\|. \quad (3.63)$$

Taking limit  $k \rightarrow \infty$  in above equation , we get

$$\lim_{k \rightarrow \infty} \|\psi_{n_k} - v_{n_k}\| = 0. \quad (3.64)$$

Similarly as in Case 1, we can prove

$$\lim_{k \rightarrow \infty} \|\kappa_{n_k} - \psi_{n_k}\| = 0 \text{ and } \lim_{k \rightarrow \infty} \|\kappa_{n_k} - \zeta_{n_k}\| = 0. \quad (3.65)$$

Also,

$$\lim_{k \rightarrow \infty} \|\zeta_{n_k} - v_{n_k}\| = 0 \text{ and } \lim_{k \rightarrow \infty} \|\zeta_{n_k+1} - \kappa_{n_k}\| = 0. \quad (3.66)$$

Similar to Case 1,  $q \in VI(L, Q)$  and  $q \in (A_j + B_j)^{-1}(0)$ . Hence,  $q \in \Omega$ . Using the same arguments as in Case 1, we have

$$\limsup_{k \rightarrow \infty} \langle g(\rho_{n_k}) - \iota, \rho_{n_k} - \iota + \gamma_{n_k}(g(\rho_{n_k}) - \rho_{n_k}) \rangle \leq 0. \quad (3.67)$$

Using equation (3.57), we obtain

$$r_{n_k+1} \leq (1 - \gamma_{n_k})r_{n_k} + \gamma_{n_k}s_{n_k}. \quad (3.68)$$

Using equation (3.59) in equation (3.68), we get

$$\gamma_{n_k}r_{n_k} \leq \gamma_{n_k}s_{n_k}. \quad (3.69)$$

As  $\gamma_{n_k} > 0$ , we have  $r_{n_k} \leq s_{n_k}$ . So, we have

$$\begin{aligned} \|\zeta_{n_k} - \iota\|^2 &\leq 2\langle g(\rho_{n_k}) - \iota, \rho_{n_k} - \iota + \gamma_{n_k}(g(\rho_{n_k}) - \rho_{n_k}) \rangle + \frac{\epsilon_{n_k}^2 \gamma_{n_k}}{\gamma_{n_k}^2} \|\zeta_{n_k} - \zeta_{n_k-1}\|^2 \\ &+ \frac{2\epsilon_{n_k}}{\gamma_{n_k}} \|\zeta_{n_k} - \iota\| \|\zeta_{n_k} - \zeta_{n_k-1}\|. \end{aligned} \quad (3.70)$$

As  $\lim_{k \rightarrow \infty} \gamma_{n_k} = 0$ ,  $\lim_{k \rightarrow \infty} \frac{\epsilon_{n_k}}{\gamma_{n_k}} \|\zeta_{n_k} - \zeta_{n_k-1}\| = 0$  and from equation (3.67), we obtain  $r_{n_k} \rightarrow 0$  as  $k \rightarrow \infty$ . Also, using equation (3.68), we have  $\lim_{k \rightarrow \infty} r_{n_k+1} = 0$  and from equation (3.60), we get  $r_{n_k+1} \geq r_k$ , which gives  $\lim_{k \rightarrow \infty} r_k = 0$  that is,  $\|\zeta_n - \iota\| \rightarrow 0$  as  $n \rightarrow \infty$ . Hence  $\zeta_n \rightarrow \iota$  as  $n \rightarrow \infty$ .  $\square$

## 4 Numerical Example

In this section, we give numerical example and compare the convergence of algorithms [23, 24] with the Algorithm (3.1).

**Example 1.** Let  $H = \mathbb{R}^4$  and  $Q = \{\zeta \in \mathbb{R}^4 : \zeta_1 + \zeta_2 - 3\zeta_3 + \zeta_4 \leq 0\}$ , where  $\zeta = (\zeta_1, \zeta_2, \zeta_3, \zeta_4)$ . Suppose  $A_j : H \rightarrow 2^H$  are maximal monotone mappings defined by  $A_j(\zeta) = N_j^* N_j(\zeta)$  for all  $\zeta \in H$ ,  $j = 1, 2$ , where  $N_j : \mathbb{R}^4 \rightarrow \mathbb{R}^4$  are created from a normal distribution with unit variance and mean zero and the mappings  $B_1, B_2 : H \rightarrow H$  are defined by  $B_1(\zeta) = \frac{\zeta}{5}$  for all  $\zeta \in H$ ,  $B_2(\zeta) = \frac{\zeta}{7}$  for all  $\zeta \in H$ . Clearly,  $B_1$  and  $B_2$  are 1- inverse strongly monotone mappings. The mappings  $S_i : H \rightarrow H$  are defined as  $S_i(\zeta) = \frac{-3i\zeta}{i+1}$  for all  $\zeta \in H, i = 1, 2, \dots, 7$ . It can be easily shown that  $S_i$  are  $\frac{2i-1}{4i+1}$  demicontractive mappings. Further,  $L : Q \rightarrow H$  is a mapping given by  $L(\zeta) = \zeta$  for all  $\zeta \in Q$ , where  $L$  is 1- cocoercive mapping and the mapping  $g : H \rightarrow H$  is defined as  $g(\zeta) = \frac{\zeta}{2}$  for all  $\zeta \in H$ . Let  $\alpha_n = \frac{1}{n^5}, \gamma_n = \frac{1}{n+6}, \epsilon = 0.4, \beta_{n,0} = \frac{57k+3}{67k}, \beta_{n,i} = \frac{10k-3}{469k}$  for all  $i = 1, 2, \dots, 7, \tau_1 = 0.5$  and  $\tau_2 = 1$  and we choose  $\|\zeta_n - \zeta_{n-1}\| \leq 10^{-4}$  as stopping criterion.

We take following cases for various initial values of  $\zeta_0$  and  $\zeta_1$  and plot the graphs of errors  $E_n = \|\zeta_n - \zeta_{n-1}\|$  against number of iterations  $n$ .

**Case 1.**  $\zeta_0 = (1, 1, 1, 1)$ ,  $\zeta_1 = (2, 2, 2, 2)$ ;

**Case 2.**  $\zeta_0 = (10, 10, 10, 10)$ ,  $\zeta_1 = (20, 20, 20, 20)$ ;

**Case 3.**  $\zeta_0 = (100, 100, 100, 100)$ ,  $\zeta_1 = (200, 200, 200, 200)$ .

We also show that Algorithm (3.1) is more effective than Lorenz algorithm [23] and Cholamjiak algorithm [24].

Cases	iteration number	cpu time in seconds
Case 1.	7	0.02158
Case 2.	10	0.02419
Case 3.	38	0.03042

Table 1: Numerical analysis of Algorithm (3.1) for various cases




Figure 1: Numerical study of Algorithm (3.1) for various values of  $\zeta_0$  and  $\zeta_1$

Algorithm	iteration number	cpu time (in seconds)
Algorithm (3.1)	7	0.01726
Lorenz Algorithm	38	0.01849
Cholamjiak Algorithm	431	0.01854

Table 2: Comparison of Algorithm (3.1) with Lorenz Algorithm [23] and Cholamjiak Algorithm [24]




Figure 2: Comparison of Algorithm (3.1) with Lorenz Algorithm [23] and Cholamjiak Algorithm [24]

## Acknowledgments

The first and fourth authors delightedly acknowledge the University Grants Commission (UGC), New Delhi for providing financial support.

## Conflict of Interest

The authors declare that there is no conflict of interest.

## References

- [1] P. L. Combettes and V. R. Wajs, Signal recovery by proximal forward–backward splitting, *Multiscale Model. Simul.*, **4** (2005), 1168–1200.
- [2] D. R. Sahu, J. C. Yao and M. Verma, Convergence rate analysis of proximal gradient methods with applications to composite minimization problems, *Optimization*, **70** (2021), 75–100.
- [3] X. Qin , N. T. An, Smoothing algorithms for computing the projection onto a Minkowski sum of convex sets, *Comput. Optim. Appl.*, **74** (2019), 821–850.
- [4] T. H. Cuong, J.C. Yao and N. D. Yen, Qualitative properties of the minimum sum-of-squares clustering problem, *Optimization*, **69** (2020), 2131–2154.



- 
- [5] Y. Wang and H. Zhang, Strong convergence of the viscosity Douglas-Rachford algorithm for inclusion problems, *Appl Set-Valued Anal. Optim.*, **2** (2020), 339–349.
- [6] R. Chugh and C. Batra, Fixed point theorems of enriched Ciric's type and enriched Hardy-Rogers contractions, *Numerical Algebra, Control and Optimization*.
- [7] X. Qin and J. C. Yao, A viscosity iterative method for a split feasibility problem, *J. Nonlinear Convex Anal.*, **20** (2019), 1497–1506.
- [8] S. Mehra, R. Chugh, S. Haque and N. Mlaiki, Iterative algorithm for solving monotone inclusion and fixed point problem of a finite family of demimetric mappings, *AIMS Mathematics*, **8** (2023), 19334-19352 .
- [9] B. Tan, S. Xu and S. Li, Inertial shrinking projection algorithms for solving hierarchical variational inequality problems, *J. Nonlinear Convex Anal.*, **21** (2020), 871–884.
- [10] Q. H. Ansari, M. Islam and J. C. Yao, Nonsmooth variational inequalities on Hadamard manifolds, *Appl Anal.*, **99** (2020), 340–358.
- [11] Y. Yao, M. Postolache and J.C. Yao, An approximation algorithm for solving a split problem of fixed point and variational inclusion, *Optimization*, (2023), 1-14.
- [12] Y. Yu, Y. Zhao and T.C. Yin, Convergence of extragradient-type methods for fixed point problems and quasimonotone variational inequalities, *J. Nonlinear Convex Anal.*, **24** (2023), 2225-2237.
- [13] L. J. Zhu, J. C. Yao and Y. Yao, Approximating solutions of a split fixed point problem of demicontractive operators, *Carpathian J. Math.*, **40** (2024), 195–206.
- [14] R. T. Rockafellar, Monotone operators and the proximal point algorithm, *SIAM J. Control Optim.*, **14** (1976), 877-898.
- [15] R. E. Bruck and S. Reich, Nonexpansive projections and resolvents of accretive operators in Banach spaces, *Houst. J. Math.*, **3** (1977), 459-470.
- [16] P. L. Lions and B. Mercier, Splitting algorithms for the sum of two nonlinear operators, *SIAM J. Numer. Anal.*, **16** (1979), 964-979.
- [17] Passty, G.B.: Ergodic convergence to a zero of the sum of monotone operators in Hilbert space. *J. Math. Anal. Appl.* **72**, 383-390 (1979).
- [18] J. Douglas and H. H. Rachford, On the numerical solution of heat conduction problems in two and three space variables, *Trans Am Math Soc.*, **82**(1956), 421–439.
- [19] D. W. Peaceman and H. H. Jr. Rachford, The numerical solution of parabolic and elliptic differential equations, *J. Appl. Ind. Math.*, **3** (1955), 28–41.
- [20] P. Tseng, A modified forward-backward splitting method for maximal monotone mappings, *SIAM J. Control Optim.*, **38** (2000), 431-446.
- [21] A. Gibali and D. V. Thong, Tseng type methods for solving inclusion problems and its applications, *Calcolo*, **55** (2018), 1-22.

- [22] B. T. Polyak, Some methods of speeding up the convergence of iteration methods, *Comput. Math. Math. Phys.*, **4** (1964), 1-17.
- [23] D. A. Lorenz and T. Pock, An inertial forward-backward algorithm for monotone inclusions, *J. Math Imaging Vis.*, **51** (2015), 311-325.
- [24] W. Cholamjiak, P. Cholamjiak and S. Suantai, An inertial forward-backward splitting method for solving inclusion problems in Hilbert spaces, *J. Fixed Point Theory Appl.*, **20** (2018), 1-17.
- [25] D. V. Thong and P. Cholamjiak, Strong convergence of a forward-backward splitting method with a new step size for solving monotone inclusions, *Comput. Appl. Math.*, **38** (2019), 1-16.
- [26] H. A. Abass, K. O. Aremu, L. O. Jolaoso and O. T. Mewomo, An inertial forward-backward splitting method for approximating solutions of certain optimization problem, *J. Nonlinear Funct. Anal.*, **2020** (2020), 1-20.
- [27] H. A. Abass, c. Izuchukwu and K. O. Aremu, A common solution of family of minimization and fixed point problem for multi-valued type-one demicontractive-type mappings, *Adv. Nonlinear Var. Inequal.*, **21** (2018), 94-108.
- [28] H. A. Abass, C. C. Okeke and O. T. Mewomo, On split equality mixed equilibrium and fixed point problems of generalized ki-strictly pseudo-contractive multivalued mappings, *Dyn. Contin. Discrete Impuls. Syst. Ser. B Appl. Algorithms*, **25** (2018), 369-395.
- [29] F. Akutsah, H. A. Abass, A. A. Mebawondu and O. K. Narain, On split generalized mixed equilibrium and fixed point problems of an infinite family of quasi-nonexpansive multivalued mappings in real Hilbert spaces, *Asian Eur. J. Math.* **15** (2021), 1-20.
- [30] P. N. Anh, H. T. C. Thach and J. K. Kim, Proximal-like subgradient methods for solving multi-valued variational inequalities, *Nonlinear Funct. Anal. Appl.*, **25** (2020), 437-451.
- [31] M. Eslamian and A. Kamandi, A novel algorithm for approximating common solution of a system of monotone inclusion problems and common fixed point problem, *J. Ind. Manag. Optim.*, **19** (2021), 1-22.
- [32] M. A. Olona and O. K. Narain, Iterative method for solving finite families of variational inequality and fixed point problems of certain multi-valued mappings, *Nonlinear Funct. Anal. Appl.*, **27** (2022), 149-167.
- [33] P. Peeyada, R. Suparatulatorn, W. Cholamjiak, An inertial Mann forward-backward splitting algorithm of variational inclusion problems and its applications, *Chaos, Solitons and Fractals* **158** (2022), 1-7.
- [34] H. K. Xu, Iterative algorithms for nonlinear operators, *J. London Math. Soc.*, **66** (2002), 240-256.
- [35] K. Afassinou, O.K. Narain and O. E. Otunuga, Iterative algorithm for approximating solutions of split monotone variational inclusion, variational inequality and fixed point problems in real Hilbert spaces, *Nonlinear Funct. Anal. Appl.*, **25** (2020), 491-510.
- [36] B. Lemaire, Which fixed point does the iteration method select? In: *Recent Advances in Optimization*, Springer, Berlin, Heidelberg, 1997.

- 
- [37] C. E. Chidume, Geometric properties of Banach spaces and nonlinear spaces and nonlinear iterations Lecture Notes in Mathematics, Springer, London, 2009.
- [38] B. Tan and S. Y. Cho, Strong convergence of inertial forward–backward methods for solving monotone inclusions, *Appl. Anal.* **101**(2021), 1-29.
- [39] C. E. Chidume and J. N. Ezeora, Krasnoselskii-type algorithm for family of multi-valued strictly pseudo-contractive mappings, *J. Fixed Point Theory Appl.*, **2014** (2014), 1-7.
- [40] P. E. Maingé, A hybrid extragradient-viscosity method for monotone operators and fixed point problems, *SIAM J Control Optim.* **47** (2008), 1499-1515.

**Anjali** Department of Mathematics, Maharshi Dayanand University, Rohtak 124001, India  
E-mail: [anjaliahuja3108@gmail.com](mailto:anjaliahuja3108@gmail.com)

**Seema Mehra** Department of Mathematics, Maharshi Dayanand University, Rohtak 124001, India  
E-mail: [sberwal2007.math@mdurohtak.ac.in](mailto:sberwal2007.math@mdurohtak.ac.in)

**Renu Chugh** Department of Mathematics, Gurugram University, Haryana 122413, India  
E-mail: [chughrenu1988@gmail.com](mailto:chughrenu1988@gmail.com)

**Charu Batra** Department of Mathematics, Maharshi Dayanand University, Rohtak 124001, India  
E-mail: [charubatra45@gmail.com](mailto:charubatra45@gmail.com)