



Existence theory for a fractional q -integral equations

Hamid Reza Sahebi and Manochehr Kazemi

Abstract. The paper focuses on establishing sufficient conditions for the existence of the solutions for a functional equation involving q -fractional integrals, particularly in Banach spaces. In this method, the technique of measures of noncompactness and Petryshyn's fixed point theorem Banach space is used. We provide some examples of equations, which confirm that our result is applicable to a wide class of integral equations.

Keywords. Measure of noncompactness, existence of solution, q -integral equation, Petryshyn's fixed point theorem

1 Introduction

The existence of solutions for integral equations (IEs) is a major challenge in mathematical research, leading to extensive investigation and development of various methodologies. Two notable frameworks for addressing solution existence are Schauder's theorem and Darbo's Fixed Point Theorem (F.P.T). A key concept in these approaches is the measure of noncompactness (M.N.C), which plays a foundational role in proving the existence results. The M.N.C is a highly significant and practical concept in non-linear analysis. It originated from Kuratowski's seminal paper [22], providing powerful tools for solving a wide range of IEs and fractional differential equations (FDEs).

In 1955, Darbo's fundamental F.P.T in [9], introduced the concept of M.N.C, which has since been widely employed by researchers to analyze the solvability of various types of IEs and DEs [1]. In fact, researchers have successfully generalized this theorem to address specific problem types they encounter. In [6], via a measure of non-compactness concept, Banas obtained an existence result nonlinear functional IEs (for more details, see [7, 29, 31]). Caballero *et al.* in [8] studied the solvability of the functional IE as follows,

$$x(t) = f\left(t, \int_0^t v(t, s, x(s)) ds, x(\alpha(t))\right) \cdot g\left(t, \int_0^a x(t)u(t, s, x(s)) ds, x(\beta(t))\right), \quad (1.1)$$

for $x \in C(I_a)$, $I_a := [0, a]$. They showed under the some conditions the functional IEs (1.1) has at least one solution $x \in C(I_a)$. In [10], authors with the help of generalized Darbo's F.P.T

established the existence of solution of a \mathbb{IE} with generalized fractional integral of two variables. Metwalli *et al.* in [24], studied the existence and the uniqueness of a.e. monotonic solutions of following problem,

$$\begin{aligned} x(t) &= h(t) + m(t) \cdot g(t, x(t - \tau)) + \int k(t, s) f(s, x(s - \tau)) ds, \\ x(t) &= \phi(t), \quad t \in [-\tau, 0]. \end{aligned} \quad (1.2)$$

Mishra *et al.* in [25] obtained some results on the existence of solutions for a nonlinear Erdelyi-Kober fractional quadratic \mathbb{IE} with deviating arguments. The solutions of system of functional \mathbb{IE} s in the setting of M.N.C investigated [26]. In [33], authors investigated existence of solutions in Banach algebra the nonlinear functional \mathbb{IE} comprising of Hadamard fractional operators under certain relevant assumptions in conjunction with F.P. theory. Using the techniques of Darbo's F.P.T associated with M.N.C, authors established the existence results for functional stochastic \mathbb{IE} s in Banach Algebra [11]. Srivastava *et al.* [34] established the existence of the solution of a functional \mathbb{IE} of two variables, which is of the form of the product of two operators in the Banach algebra $C(I_a \times I_a)$.

The existence of solution of q -integral equations (q - \mathbb{IE} s) have been analyzed by several researchers [2, 4, 5, 23, 30, 32]. In [17], the authors discussed the q - \mathbb{IE} ,

$$x(t) = F\left(t, x(a(t)), \frac{x(b(t))}{\Gamma_q(\alpha)} \int_0^t (t - qs)^{(\alpha-1)} u(s, x(s)) d_qs\right), \quad t \in I_1, \quad (1.3)$$

where $\alpha > 1$, $q \in (0, 1)$. With a generalized version of Darbo's theorem, they have established a sufficient conditions for the existence of at least one solution to (1.3). In [3], the author examined quadratic \mathbb{IE} as follows,

$$y(t) = y(t) + \frac{(Ay)(t)}{\Gamma_q(\beta)} \int_0^t k(t, s)(t - qs)^{(\beta-1)} (By)(s) d_qs, \quad t \in I_1, \quad (1.4)$$

where $\beta > 0$, $q \in (0, 1)$ and $A, B : C(I_1) \rightarrow C(I_1)$. Recently, Kazemi and Ezzati [19] utilized Petryshyn's F.P.T for the existence of solutions of nonlinear functional \mathbb{IE} s. They showed that Petryshyn's theorem is much more useful than Darbo's theorem. In fact, it dose not require to confirm the used operator of maps a closed convex subset onto itself [20].

Motivated and stimulated by the [19, 20], we introduced and analyze the existence result for the following a class of functional q - \mathbb{IE} with fractional order,

$$\varkappa(t) = \Psi\left(t, \varkappa(a(t)), \pi(t, \varkappa(c(t))), \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} p(t, s, \varkappa(b(s))) d_qs\right), \quad (1.5)$$

where $0 < q < 1$, $\beta > 0$ and $\varsigma > 1$.

The paper is structured as follows. In Section 2, we collect some definitions, lemmas and theorems, which are essential to prove our main results. In Section 3, we introduce a new functional q - \mathbb{IE} for existence of a solution a Banach space. Then, we establish and prove a existence theorem by utilized Petryshyn's F.P.T. In Section 4, we also give some examples to support our main theorem. Finally, in Section 5, concludes the paper.

2 Auxiliary facts and notations

In this section, we review some definitions and theorems, by stating some auxiliary facts and notations. Let

- X : Banach space,
- N_ε : A ball of radius ε ,
- \bar{N}_ε : Sphere in X with radius ε ,
- $C(I_a)$: All real functions continuous on I_a ,
- $B(X)$: Bounded set of X ,
- $\text{Fix}(T)$: the set of fixed points of T in X .

In 1910, Jackson introduced the concept of quantum calculus in [16]. There are several applications of q -calculus in physical problems such as molecular problems [13], elementary particle physics, and chemical physics [12, 15]. Basic definitions and properties of quantum calculus can be found in the book [18]. In follow, we recall some basic facts on quantum calculus and present additional properties that will be used later.

For $q \in (0, 1)$, we define $[\varkappa]_q = \frac{1-q^\varkappa}{1-q}$, $\varkappa \in \mathbb{R}$. The q -analogue of the power function $(1 - \vartheta)^k$ with $k \in \{0, 1, 2, \dots\}$ is,

$$(1 - \vartheta)^{(0)} = 1, \quad (1 - \vartheta)^{(k)} = \prod_{i=0}^{k-1} (1 - \vartheta q^i), \quad k \in \mathbb{N}, \vartheta \in \mathbb{R}. \quad (2.1)$$

Also, The q -integral of a function f defined on the interval I_a is given by,

$$\int_0^\nu f(\zeta) d_q \zeta = \nu(1 - q) \sum_{i=0}^{\infty} f(\nu q^i) q^i, \quad \nu \in I_a. \quad (2.2)$$

For any $m, n > 0$, $B_q(m, n) = \int_0^1 \zeta^{(m-1)} (1 - q\zeta)^{(n-1)} d_q \zeta$ is called the q -beta function.

Lemma 2.1 ([28]). *Let $m, n > 0$, and $0 < q < 1$. Then we have*

$$\int_0^\nu (\nu - q\zeta)^{(m-1)} \zeta^{(n)} d_q \zeta = \nu^{m+n} B_q(m, n + 1). \quad (2.3)$$

Definition 1 ([22]). Let $T \subset B(X)$, then

$$\phi(T) = \inf\{\varepsilon : \text{there exist a finite number of sets of diameter } \leq \varepsilon \text{ that can cover } T\}, \quad (2.4)$$

is said to be the Kuratowski M.N.C.

Definition 2 ([14]). Let $T \subset B(X)$, then

$$\zeta(T) = \inf \left\{ \varepsilon > 0 : T \text{ has a finite } \varepsilon - \text{net in } X \right\}, \quad (2.5)$$

is said to be the Hausdorff M.N.C.

Definition 3 ([14]). For $T \subset B(X)$, the M.N.C ϕ and ζ fulfill $\zeta(T) \leq \phi(T) \leq 2\zeta(T)$.

For $h \in C(I_a)$ and $\varepsilon \geq 0$ denoted by $\delta_\varepsilon(h)$, the modulus of continuity of the function h , i.e.,

$$\delta_\varepsilon(h) = \sup \left\{ |h(t) - h(\hat{t})| : |t - \hat{t}| \leq \varepsilon \right\}. \quad (2.6)$$

The uniformly continuous h on I_a implies that $\delta_\varepsilon(h) \rightarrow 0$ as $\varepsilon \rightarrow 0$.

Theorem 2.1 ([21]). *For $T \subset B(X)$, the M.N.C in $C(I_a)$ is denoted by,*

$$T = \limsup_{\varepsilon \rightarrow 0} \delta_\varepsilon t, \quad t \in T. \quad (2.7)$$

Theorem 2.2 ([27]). *Let $\mathfrak{F}, \mathfrak{E} \subset X$ then*

1. $\zeta(\mathfrak{F} \cup \mathfrak{E}) = \max \{ \zeta(\mathfrak{F}), \zeta(\mathfrak{E}) \};$
2. $\zeta(\mathfrak{F} + \mathfrak{E}) \leq \zeta(\mathfrak{F}) + \zeta(\mathfrak{E});$
3. $\zeta(\varsigma \mathfrak{F}) = |\varsigma| \zeta(\mathfrak{F})$, where $\varsigma \mathfrak{F} = \{ \varsigma m : m \in \mathfrak{F} \};$
4. $\zeta(\mathfrak{F}) \leq \zeta(\mathfrak{E})$, for $\mathfrak{F} \subset \mathfrak{E};$
5. $\zeta(\bar{c} \circ \mathfrak{F}) = \zeta(\mathfrak{F});$

Recall that $F \in C(X)$ is called a κ -set contraction if for $K \subset B(X)$, F is bounded and $\phi(FK) \leq \kappa \phi(K)$ for each $0 < \kappa < 1$. Moreover, If $\phi(FK) \leq \phi(K)$, F is called condensing map.

Let us recall that Petryshyn's F.P.T,

Theorem 2.3 ([27]). *Let $F : N_\varepsilon \rightarrow X$ be a condensing mapping and satisfying the following boundary condition,*

$$\text{if } F(x) = \kappa x, \text{ for some } x \in \bar{N}_\varepsilon, \text{ with } \kappa \leq 1. \quad (2.8)$$

Then $\text{Fix}_{N_\varepsilon}(F) \neq \emptyset$.

In order to prove the main results, we first establish the following lemma.

Lemma 2.2. *The following inequality is hold*

$$\begin{aligned} & \left| \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} p(t, s, \varkappa(b(s))) d_qs \right. \\ & \quad \left. - \int_0^{r(\hat{t})} (r(\hat{t}) - qs)^{(\varsigma-1)} p(\hat{t}, s, \varkappa(b(s))) d_qs \right| \\ & \leq \hat{\delta}_\tau(u) D^\varsigma B_q(\varsigma, 1) + 2LB_q(\varsigma, 1) \delta_\tau(w)^\varsigma, \end{aligned} \quad (2.9)$$

where

$$\begin{aligned} \delta_\tau(r) &= \sup \left\{ |r(t) - r(\hat{t})| : t, \hat{t} \in I_a, |t - \hat{t}| \leq \tau \right\}, \\ L &= \sup \left\{ |p(t, s, z)| : t \in I_a, s \in [0, D], z \in [-\varepsilon, \varepsilon] \right\}, \\ \hat{\delta}_\tau(u) &= \sup \left\{ |p(t, x, y) - p(\hat{t}, x, y)| : |t - \hat{t}| \leq \tau, x \in [0, D], y \in [-\varepsilon, \varepsilon] \right\}. \end{aligned} \quad (2.10)$$

Proof. We have

$$\begin{aligned}
& \left| \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} p(t, s, \varkappa(b(s))) \, d_qs \right. \\
& \quad \left. - \int_0^{r(\hat{t})} (r(\hat{t}) - qs)^{(\varsigma-1)} p(\hat{t}, s, \varkappa(b(s))) \, d_qs \right| \\
& \leq \left| \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} p(t, s, \varkappa(b(s))) \, d_qs \right. \\
& \quad \left. - \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} p(\hat{t}, s, \varkappa(b(s))) \, d_qs \right| \\
& \quad + \left| \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} p(\hat{t}, s, \varkappa(b(s))) \, d_qs \right. \\
& \quad \left. - \int_0^{r(\hat{t})} (r(t) - qs)^{(\varsigma-1)} p(\hat{t}, s, \varkappa(b(s))) \, d_qs \right| \\
& \quad + \left| \int_0^{r(\hat{t})} (r(t) - qs)^{(\varsigma-1)} p(\hat{t}, s, \varkappa(b(s))) \, d_qs \right. \\
& \quad \left. - \int_0^{r(\hat{t})} (r(\hat{t}) - qs)^{(\varsigma-1)} p(\hat{t}, s, \varkappa(b(s))) \, d_qs \right| \\
& \leq \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} \left| p(t, s, \varkappa(b(s))) - p(\hat{t}, s, \varkappa(b(s))) \right| \, d_qs \\
& \quad + \int_{r(\hat{t})}^{r(t)} (r(t) - qs)^{(\varsigma-1)} \left| p(\hat{t}, s, \varkappa(b(s))) \right| \, d_qs \\
& \quad + \int_0^{r(\hat{t})} [(r(\hat{t}) - qs)^{(\varsigma-1)} - (r(t) - qs)^{(\varsigma-1)}] \left| p(\hat{t}, s, \varkappa(b(s))) \right| \, d_qs \\
& \leq \hat{\delta}_\tau(u) \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} \, d_qs \\
& \quad + L \int_0^{r(\hat{t})} (r(\hat{t}) - qs)^{(\varsigma-1)} - (r(t) - qs)^{(\varsigma-1)} \, d_qs \\
& \quad + \int_{r(\hat{t})}^{r(t)} (r(t) - qs)^{(\varsigma-1)} \, d_qs \\
& = \hat{\delta}_\tau(u) r(t)^\varsigma B_q(\varsigma, 1) + L B_q(\varsigma, 1) \left[r(\hat{t})^\varsigma - r(t)^\varsigma + 2(r(t) - r(\hat{t}))^\varsigma \right] \\
& \leq \hat{\delta}_\tau(u) r(t)^\varsigma B_q(\varsigma, 1) + 2L B_q(\varsigma, 1) (r(t) - r(\hat{t}))^\varsigma \\
& \leq \hat{\delta}_\tau(u) D^\varsigma B_q(\varsigma, +1) + 2L B_q(\varsigma, 1) \delta(r, \tau)^\varsigma.
\end{aligned} \tag{2.11}$$

□

3 A Existence Theorem

In this section, we will study the existence of q - \mathbb{IE} (1.5) under the following assumptions,

- S1) $\varkappa \in C(I_a)$, $\Psi \in C(I_a \times \mathbb{R}^{\mathbb{K}})$, $\pi \in C(I_a \times \mathbb{R})$, $p \in C(I_a \times [0, D] \times \mathbb{R})$, $c : I_a \rightarrow I_a$, $b : \mathbb{R}^+ \rightarrow I_a$, $r : I_a \rightarrow \mathbb{R}^+$ are continuous and for each $t \in I_a$ and $r(t) \leq D$;
- S2) There exist nonnegative constant k_1, k_2, k_3 and k_4 with $k_1 + k_2 k_4 \leq 1$ s.t.

$$|\Psi(t, t_1, t_2, t_3) - \Psi(t, \hat{t}_1, \hat{t}_2, \hat{t}_3)| \leq k_1 |t_1 - \hat{t}_1| + k_2 |t_2 - \hat{t}_2| + k_3 |t_3 - \hat{t}_3|, \quad (3.1)$$

$$\text{and } |\pi(x, y) - \pi(x, \hat{y})| \leq k_4 |y - \hat{y}|;$$

- S3) For $\varepsilon \geq 0$, the operator Ψ satisfies the following condition,

$$\sup \left\{ |\Psi(t, x, y, z)| : t \in I_a, x, y \in [-\varepsilon, \varepsilon], \right. \\ \left. - \frac{D^\varsigma}{\Gamma_q(\varsigma)} L B_q(\varsigma, 1) \leq z \leq \frac{D^\varsigma}{\Gamma_q(\varsigma)} L B_q(\varsigma, 1) \right\} \leq \varepsilon, \quad (3.2)$$

$$\text{where } L = \sup \left\{ |p(t, s, z)| : t \in I_a, s \in [0, D], z \in [-\varepsilon, \varepsilon] \right\}.$$

Theorem 3.1. *Under the assumption (S1)-(S3), the functional q - \mathbb{IE} (1.5) has at least one solution in $C(I_a)$.*

Proof. Let

$$\left\{ \begin{array}{l} F : N_\varepsilon \rightarrow X, \\ (F \varkappa)(t) = \Psi \left(t, \varkappa(a(t)), \pi(t, \varkappa(c(t))), \right. \\ \left. \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} p(t, s, \varkappa(b(s))) \, d_q s \right). \end{array} \right. \quad (3.3)$$

We divided the proof into several steps.

Step 1. The operator F is continuous on the ball N_ε . Consider $\tau > 0$ and any $\varkappa, \hat{\varkappa} \in N_\varepsilon$ s.t. $\|\varkappa - \hat{\varkappa}\| < \tau$. We have,

$$\begin{aligned} |(F \varkappa)(t) - (F \hat{\varkappa})(t)| &= \left| \Psi \left(t, \varkappa(a(t)), \pi(t, \varkappa(c(t))), \right. \right. \\ &\quad \left. \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} p(t, s, \varkappa(b(s))) \, d_q s \right) \\ &\quad \left. - \Psi \left(t, \hat{\varkappa}(a(t)), \pi(t, \hat{\varkappa}(c(t))), \right. \right. \\ &\quad \left. \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} p(t, s, \hat{\varkappa}(b(s))) \, d_q s \right) \Big| \\ &\leq \left| \Psi \left(t, \varkappa(a(t)), \pi(t, \varkappa(c(t))), \right. \right. \\ &\quad \left. \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} p(t, s, \varkappa(b(s))) \, d_q s \right) \\ &\quad \left. - \Psi \left(t, \hat{\varkappa}(a(t)), \pi(t, \varkappa(c(t))), \right. \right. \\ &\quad \left. \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} p(t, s, \varkappa(b(s))) \, d_q s \right) \Big| \end{aligned}$$

$$\begin{aligned}
& + \left| \Psi \left(t, \hat{\varkappa}(a(t)), \pi(t, \varkappa(c(t))), \right. \right. \\
& \quad \left. \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} p(t, s, \varkappa(b(s))) d_qs \right) \\
& \quad - \Psi \left(t, \hat{\varkappa}(a(t)), \pi(t, \hat{\varkappa}(c(t))), \right. \\
& \quad \left. \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} p(t, s, \varkappa(b(s))) d_qs \right) \Big| \\
& + \left| \Psi \left(t, \hat{\varkappa}(a(t)), \pi(t, \hat{\varkappa}(c(t))), \right. \right. \\
& \quad \left. \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} p(t, s, \varkappa(b(s))) d_qs \right) \\
& \quad - \Psi \left(t, \hat{\varkappa}(a(t)), \pi(t, \hat{\varkappa}(c(t))), \right. \\
& \quad \left. \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} p(t, s, \hat{\varkappa}(b(s))) d_qs \right) \Big| \\
& \leq k_1 \left| \varkappa(a(t)) - \hat{\varkappa}(a(t)) \right| + k_2 \left| g(t, \varkappa(c(t))) - g(t, \hat{\varkappa}(c(t))) \right| \\
& \quad + \frac{k_3}{\Gamma_q(\varsigma)} \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} \left| p(t, s, \varkappa(b(s))) - p(t, s, \hat{\varkappa}(b(s))) \right| d_qs \\
& \leq k_1 \left| \varkappa(a(t)) - \hat{\varkappa}(a(t)) \right| + k_2 k_4 \left| \varkappa(c(t)) - \hat{\varkappa}(c(t)) \right| \\
& \quad + \frac{k_3}{\Gamma_q(\varsigma)} \delta_\tau(u) r(t)^{(\varsigma)} B_q(\varsigma, 1) \\
& \leq (k_1 + k_2 k_4) \|\varkappa - \hat{\varkappa}\| + \frac{k_3}{\Gamma_q(\varsigma)} \delta_\tau(u) D^\varsigma B_q(\varsigma, 1), \tag{3.4}
\end{aligned}$$

where

$$\begin{aligned}
\delta_\tau(u) = \sup \Big\{ & |p(t, x, y) - p(t, x, \hat{y})| : t \in I_a, x \in [0, D], \\
& y, \hat{y} \in [-\varepsilon, \varepsilon], |y - \hat{y}| \leq \varepsilon \Big\}. \tag{3.5}
\end{aligned}$$

The uniform continuity of $p(\cdot, \cdot, \cdot)$ on the subset $I_a \times [0, D] \times \mathbb{R}$ implies that $\delta_\tau(u) \rightarrow 0$ as $\tau \rightarrow 0$. This shows that the operator F is continuous on N_ε .

Step 2. The densifying condition operator F . Let E_τ be a bounded subset in X and $t, \hat{t} \in I_a$ such that $t \leq \hat{t}$ and $t - \hat{t} \leq \tau$ for $\tau > 0$, Lemma 2.2 implies that,

$$\begin{aligned}
|(F\varkappa)(t) - (F\varkappa)(\hat{t})| & = \left| \Psi \left(t, \varkappa(a(t)), \pi(t, \varkappa(c(t))), \right. \right. \\
& \quad \left. \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} p(t, s, \varkappa(b(s))) d_qs \right) \\
& \quad - \Psi \left(\hat{t}, \varkappa(a(\hat{t})), \pi(\hat{t}, \varkappa(c(\hat{t}))), \right. \\
& \quad \left. \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(\hat{t})} (r(\hat{t}) - qs)^{(\varsigma-1)} p(\hat{t}, s, \varkappa(b(s))) d_qs \right) \Big|
\end{aligned}$$

$$\begin{aligned}
&\leq \left| \Psi \left(t, \varkappa(a(t)), \pi(t, \varkappa(c(t))), \right. \right. \\
&\quad \left. \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} p(t, s, \varkappa(b(s))) \, d_qs \right) \\
&\quad \left. - \Psi \left(t, \varkappa(a(t)), \pi(t, \varkappa(c(t))), \right. \right. \\
&\quad \left. \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(\hat{t})} (r(\hat{t}) - qs)^{(\varsigma-1)} p(\hat{t}, s, \varkappa(b(s))) \, d_qs \right) \Big| \\
&\quad + \left| \Psi \left(t, \varkappa(a(t)), \pi(t, \varkappa(c(t))), \right. \right. \\
&\quad \left. \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(\hat{t})} (r(\hat{t}) - qs)^{(\varsigma-1)} p(\hat{t}, s, \varkappa(b(s))) \, d_qs \right) \\
&\quad \left. - \Psi \left(t, \varkappa(a(t)), \pi(\hat{t}, \varkappa(c(\hat{t}))), \right. \right. \\
&\quad \left. \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(\hat{t})} (r(\hat{t}) - qs)^{(\varsigma-1)} p(\hat{t}, s, \varkappa(b(s))) \, d_qs \right) \Big| \\
&\quad + \left| \Psi \left(t, \varkappa(a(\hat{t})), \pi(\hat{t}, \varkappa(c(\hat{t}))), \right. \right. \\
&\quad \left. \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(\hat{t})} (r(\hat{t}) - qs)^{(\varsigma-1)} p(\hat{t}, s, \varkappa(b(s))) \, d_qs \right) \Big| \\
&\quad - \Psi \left(t, \varkappa(a(\hat{t})), \pi(\hat{t}, \varkappa(c(\hat{t}))), \right. \\
&\quad \left. \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(\hat{t})} (r(\hat{t}) - qs)^{(\varsigma-1)} p(\hat{t}, s, \varkappa(b(s))) \, d_qs \right) \Big| \\
&\quad + \left| \Psi \left(t, f(\hat{t}, \varkappa(a(\hat{t}))), \pi(\hat{t}, \varkappa(c(\hat{t}))), \right. \right. \\
&\quad \left. \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(\hat{t})} (r(\hat{t}) - qs)^{(\varsigma-1)} p(\hat{t}, s, \varkappa(b(s))) \, d_qs \right) \\
&\quad \left. - \Psi \left(\hat{t}, f(\hat{t}, \varkappa(a(\hat{t}))), \pi(\hat{t}, \varkappa(c(\hat{t}))), \right. \right. \\
&\quad \left. \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(\hat{t})} (r(\hat{t}) - qs)^{(\varsigma-1)} p(\hat{t}, s, \varkappa(b(s))) \, d_qs \right) \Big| \\
&\leq \frac{k_3}{\Gamma_q(\varsigma)} \left| \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} p(t, s, \varkappa(b(s))) \, d_qs \right. \\
&\quad \left. - \int_0^{r(\hat{t})} s(r(\hat{t}) - qs)^{(\varsigma-1)} p(\hat{t}, s, \varkappa(b(s))) \, d_qs \right| \\
&\quad + k_2 \left| g(t, \varkappa(c(t))) - \pi(t, \varkappa(c(\hat{t}))) \right| + k_2 \left| g(t, \varkappa(c(t))) - \pi(\hat{t}, \varkappa(c(\hat{t}))) \right| \\
&\quad + k_1 \left| \varkappa(a(t)) - \varkappa(a(\hat{t})) \right| + k_1 \left| \varkappa(a(t)) - f(\hat{t}, \varkappa(a(\hat{t}))) \right| + \delta(\Psi, \tau)
\end{aligned}$$

$$\begin{aligned} &\leq \frac{k_3 \hat{\delta}_\tau(u)}{\Gamma_q(\varsigma)} D^\varsigma B_q(\varsigma, 1) + \frac{2k_3 L}{\Gamma_q(\varsigma)} B_q(\varsigma, 1) \delta_\tau(w)^\varsigma + k_1 \left| \varkappa(a(t)) - \varkappa(a(\hat{t})) \right| \\ &\quad + k_2 k_4 \left| \varkappa(c(t)) - \varkappa(c(\hat{t})) \right| k_1 \delta_\tau(\pi) + k_2 \delta_\tau(f) + \delta_\tau(\Psi), \end{aligned} \quad (3.6)$$

where

$$\begin{aligned} \delta_\tau(\Psi) &= \sup \left\{ |\Psi(t, x, y, z) - \Psi(\hat{t}, x, y, z)| : |t - \hat{t}| \leq \tau, \right. \\ &\quad \left. x, y \in [-\varepsilon, \varepsilon], z \in [-DL, DL] \right\}, \\ \delta_\tau(r) &= \sup \left\{ |r(t) - r(\hat{t})| : t, \hat{t} \in I_a, |t - \hat{t}| \leq \tau \right\}, \\ \delta_\tau(\pi) &= \sup \left\{ |\pi(t, x) - \pi(\hat{t}, x)| : |t - \hat{t}| \leq \tau, x \in [-\varepsilon, \varepsilon] \right\}, \\ \hat{\delta}_\tau(u) &= \sup \left\{ |p(t, x, y) - p(\hat{t}, x, y)| : |t - \hat{t}| \leq \tau, x \in [0, D], y \in [-\varepsilon, \varepsilon] \right\}. \end{aligned} \quad (3.7)$$

So

$$\begin{aligned} \delta(F\varkappa, \tau) &\leq \frac{k_3 \hat{\delta}_\tau(u)}{\Gamma_q(\varsigma)} D^\varsigma B_q(\varsigma, 1) + \frac{2k_3 L}{\Gamma_q(\varsigma)} B_q(\varsigma, 1) \delta_\tau(r)^\varsigma \\ &\quad + k_1 \delta(\varkappa, \delta_\tau(a(t))) + k_2 k_4 \delta(\varkappa, \delta_\tau(c(t))). \end{aligned} \quad (3.8)$$

Now, by taking limit of (3.8) equation as $\tau \rightarrow 0$ would give easily following inequality

$$\delta(F\varkappa, \tau) \leq (k_1 + k_2 k_4) \delta(\varkappa, \tau), \quad (3.9)$$

which this provide $\zeta(FE_\tau) \leq (k_1 + k_2 k_4) \zeta(E_\tau)$. Therefore, Ψ is condensing map.

Step 3. The boundary condition. Suppose that $z \in \tilde{N}_\varepsilon$. If $\|Fz\| = k\|z\| = k\varepsilon$, the condition (S3) implies that

$$\begin{aligned} \left| (Fz)(t) \right| &= \left| \Psi \left(t, \varkappa(a(t)), \pi(t, \varkappa(c(t))), \right. \right. \\ &\quad \left. \left. \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} p(t, s, \varkappa(b(s))) d_qs \right) \right| \leq \varepsilon, \quad \forall t \in I_a. \end{aligned} \quad (3.10)$$

Hence $\|Fz\| \leq \varepsilon$, i.e., $k \leq 1$. This complete the proof. \square

Corollary 3.2. Assume that the following conditions are holds,

(P1) $\varkappa \in C(I_a \times I_a \times \mathbb{R})$, $g \in C(I_a \times \mathbb{R})$, $r \in C(I_a, \mathbb{R}^+)$, $r(t) \leq D$, and there exist positive constant l_i , $i = 1, \dots, 5$ s.t.

$$|\Psi(t, x, y) - \Psi(t, \bar{x}, \bar{y})| \leq l_1 |x - \bar{x}| + l_2 |y - \bar{y}|, \quad (3.11)$$

$$|\Psi(t, 0, 0)| \leq l_3, \quad |\pi(t, x) - \pi(t, \bar{x})| \leq l_4 |x - \bar{x}|, \quad \text{and } |\pi(t, 0)| \leq l_5;$$

(P2) $p(t, s, x) \in C(I_a \times I_a \times \mathbb{R})$ and $\exists c_1, c_2 > 0$ s.t. $|p(t, s, x)| \leq c_1 + c_2 |x|$, $\forall t, s \in I_a, x \in \mathbb{R}$;

(P3) $l_1 + \frac{1}{\Gamma_q(\varsigma)} l_2 (l_4 + l_5) D^\varsigma B_q(\varsigma, 1) (c_1 + c_2) < 1$.

Then

$$\varkappa(t) = \Psi \left(t, \varkappa(a(t)), \frac{\pi(t, \varkappa(c(t)))}{\Gamma_q(\varsigma)} \int_0^{r(t)} (t - qs)^{(\varsigma-1)} p(t, s, \varkappa(b(s))) d_qs \right), \quad (3.12)$$

has at least one solution in $C(I_a)$.

Proof. One can easily check, (P1) implied that (S2) in Theorem 3.1. Now, we prove (S3) holds. Let $\varepsilon > 0$, $\|\varkappa\| \leq \varepsilon$. We obtain,

$$\begin{aligned}
|\varkappa(t)| &= \left| \Psi\left(t, \varkappa(a(t)), \frac{\pi(t, \varkappa(c(t)))}{\Gamma_q(\varsigma)} \int_0^{r(t)} (t - qs)^{(\varsigma-1)} p(t, s, \varkappa(b(s))) d_qs\right) \right| \\
&\leq \left| \Psi\left(t, \varkappa(a(t)), \frac{\pi(t, \varkappa(c(t)))}{\Gamma_q(\varsigma)} \int_0^{r(t)} (t - qs)^{(\varsigma-1)} p(t, s, \varkappa(b(s))) d_qs\right) \right. \\
&\quad \left. - \Psi(t, 0, 0) \right| + |\Psi(t, 0, 0)| \\
&\leq l_1 \|\varkappa(a(t))\| \\
&\quad + l_2 \left| \frac{\pi(t, \varkappa(c(t)))}{\Gamma_q(\varsigma)} \int_0^{r(t)} (t - qs)^{(\varsigma-1)} p(t, s, \varkappa(b(s))) d_qs \right| + l_3 \\
&\leq l_1 \|\varkappa(a(t))\| + l_2 \left(|\pi(t, \varkappa(c(t))) - \pi(t, 0)| \right. \\
&\quad \left. + |\pi(t, 0)| \right) \frac{1}{\Gamma_q(\varsigma)} \left| \int_0^{r(t)} (t - qs)^{(\varsigma-1)} p(t, s, \varkappa(b(s))) d_qs \right| \\
&\leq l_1 \|\varkappa(a(t))\| + l_2 \left(l_4 \|\varkappa(c(t))\| \right. \\
&\quad \left. + l_5 \right) \frac{1}{\Gamma_q(\varsigma)} r(t)^\varsigma B_q(\varsigma, 1) (c_1 + c_2 \|\varkappa(c(t))\|). \tag{3.13}
\end{aligned}$$

Hence, ε in (S3) is real number that satisfies

$$l_1 \varepsilon + \frac{1}{\Gamma_q(\varsigma)} l_2 (l_4 \varepsilon + l_5) D^\varsigma B_q(\varsigma, 1) (c_1 + c_2 \varepsilon) \leq \varepsilon. \tag{3.14}$$

Now, we define the continuous function $\Sigma : [0, 1] \rightarrow \mathbb{R}$ as follows,

$$\Sigma(\varepsilon) = l_1 \varepsilon + \frac{1}{\Gamma_q(\varsigma)} l_2 (l_4 \varepsilon + l_5) D^{\varsigma+} B_q(\varsigma, 1) (c_1 + c_2 \varepsilon) - \varepsilon. \tag{3.15}$$

The property (P3) implies that $\Sigma(0)\Sigma(1) < 0$, then $\exists \varepsilon \in (0, 1)$ s.t. $\Sigma(\varepsilon) = 0$. \square

Remark 1. Jleli *et al.* [17] introduced a functional q - \mathbb{IE} (1.3) that include the assumptions (A1)-(A9). They proved the functional q - \mathbb{IE} has at least one solution in $C([0, 1])$. By utilization of suggestion technique in Theorem 3.1 and Corollary 3.2, one can achieve the desired result with the least conditions compared to Jleli article.

Corollary 3.3. *Let*

- 1) $\varkappa \in C(I_a, \mathbb{R})$, $\varphi \in C(I_a \times \mathbb{R}^2)$, $f, \pi \in C(I_a \times \mathbb{R})$, $p \in C(I_a \times [0, D] \times \mathbb{R})$, and $c : I_a \rightarrow I_a$, $b : \mathbb{R}^+ \rightarrow I_a$, $r : I_a \rightarrow \mathbb{R}^+$ are continuous and for each $t \in I_a$, $r(t) \leq D$;
- 2) There exist nonnegative constant k_1, k_2, k_3 and k_4 with $k_1 k_3 + k_4 \leq 1$ s.t.,

$$|\varphi(t, t_1, t_2) - \Psi(t, \hat{t}_1, \hat{t}_2)| \leq k_1 |t_1 - \hat{t}_1| + k_2 |t_2 - \hat{t}_2|, \tag{3.16}$$

$$|\pi(x, y) - \pi(x, \hat{y})| \leq k_3 |y - \hat{y}|, \text{ and } |f(s, y) - f(s, \hat{y})| \leq k_4 |y - \hat{y}|;$$

3) For $\varepsilon \geq 0$, the operator Ψ satisfies the following condition,

$$\sup \left\{ |\varphi(t, y, z)| : t \in I_a, y \in [-\varepsilon, \varepsilon], \right. \\ \left. -\frac{D^\varsigma}{\Gamma_q(\varsigma)} L B_q(\varsigma, 1) \leq z \leq \frac{D^\varsigma}{\Gamma_q(\varsigma)} L B_q(\varsigma, 1) \right\} \leq \varepsilon, \quad (3.17)$$

where $L = \sup\{|p(t, s, z)| : t \in I_a, s \in [0, D], z \in [-\varepsilon, \varepsilon]\}$.

Then

$$\begin{aligned} \varkappa(t) = & f(t, \varkappa(a(t))) + \varphi\left(t, \pi(t, \varkappa(c(t))), \right. \\ & \left. \frac{1}{\Gamma_q(\varsigma)} \int_0^{r(t)} (r(t) - qs)^{(\varsigma-1)} p(t, s, \varkappa(b(s))) d_qs \right), \end{aligned} \quad (3.18)$$

has at least one solution in $C(I_b)$.

4 Examples

In this section, based on the explained approach, we incline to present some examples by using Maple software to grantee Theorem 3.1.

Example 1. Consider the following nonlinear q - \mathbb{IE} ,

$$\begin{aligned} \varkappa(t) = & \frac{1}{2} e^{-t^2} \sin \frac{\varkappa(t)}{2} + \frac{e^{-\sqrt{t}t^2}}{4+4t^2} \ln(1 + |\varkappa(t)|) \\ & + \frac{1}{\Gamma_{1/2}(\frac{3}{2})} \int_0^{\sqrt{t}} \left(\sqrt{t} - \frac{1}{2}s \right)^{1/2} \frac{\varkappa(s)}{2+s^2} d_qs, \quad t \in I_1. \end{aligned} \quad (4.1)$$

The assumptions (S1) and (S2) of Theorem 3.1 are satisfied. Now, we check that (S3) also holds. Suppose that $\|\varkappa(t)\| \leq \varepsilon$, then,

$$\begin{aligned} |\varkappa(t)| = & \left| \frac{1}{2} e^{-t^2} \sin \frac{\varkappa(t)}{2} + \frac{e^{-\sqrt{t}t^2}}{4+4t^2} \ln(1 + |\varkappa(t)|) \right. \\ & \left. + \frac{1}{\Gamma_{1/2}(\frac{3}{2})} \int_0^{\sqrt{t}} \left(\sqrt{t} - \frac{1}{2}s \right)^{1/2} \frac{\varkappa(s)}{2+s^2} d_qs \right| \\ \leq & \frac{1}{2} + \frac{1}{4}\varepsilon + \frac{1}{2\Gamma_{1/2}(\frac{3}{2})} B_{1/2} \left(\frac{3}{2}, 1 \right) \varepsilon \leq \varepsilon. \end{aligned} \quad (4.2)$$

Hence, (S3) holds if, $\varepsilon \geq 1.51496$. This implies that the equation has at least one solution in $C(I_1)$.

Example 2. Consider the following nonlinear q - \mathbb{IE} ,

$$\varkappa(t) = \frac{1}{4} (te^{-t} + t^3 \varkappa(t))$$

$$+ \frac{1}{\Gamma_{1/4}(\frac{5}{2})} \int_0^{t^2} \left(t^2 - \frac{1}{4}s\right)^{3/2} \left(\frac{1}{2}\varkappa(s^2) + se^{-2t} \frac{\sin(t)}{2 + |\cos(t)|}\right) d_qs. \quad (4.3)$$

One can see that, the assumptions (S1) and (S2) of Theorem 3.1 are satisfied. Now, we check that (S3) also holds. Suppose that $\|\varkappa(t)\| \leq \varepsilon$, then,

$$\begin{aligned} |\varkappa(t)| &= \left| \frac{1}{4} (te^{-t} + t^3 \varkappa(t)) \right. \\ &\quad \left. + \frac{1}{\Gamma_{1/4}(\frac{5}{2})} \int_0^{t^2} \left(t^2 - \frac{1}{4}s\right)^{3/2} \left(\frac{1}{2}\varkappa(s^2) + se^{-2t} \frac{\sin(t)}{2 + |\cos(t)|}\right) d_qs \right| \\ &\leq \frac{1}{4}(\varepsilon + 1) + \frac{1}{\Gamma_{1/4}(\frac{5}{2})} B_{1/4} \left(\frac{5}{2}, 1\right) \left(\frac{1}{2}\varepsilon + \frac{1}{2}\right) \leq \varepsilon. \end{aligned} \quad (4.4)$$

Thus, (S3) holds if, $\varepsilon \geq 1.50045$. This implies that the equation has at least one solution in $C(I_1)$.

Example 3. Consider the following nonlinear q - \mathbb{IE} ,

$$\begin{aligned} \varkappa(t) &= \frac{e^{-2t}}{3} + \frac{\ln(1 + |\varkappa(\sqrt{t})|)}{4 + t^2} + \frac{|\varkappa(\sqrt{t})|}{1 + |\varkappa(\sqrt{t})|} \\ &\quad + \frac{1}{\Gamma_{1/4}(\frac{5}{2})} \int_0^{t^2} \left(t^2 - \frac{1}{4}s\right)^{3/2} \sqrt{\frac{1 + \varkappa(\sqrt{t})}{1 + st}} d_qs, \quad t \in [0, 1]. \end{aligned} \quad (4.5)$$

The assumptions (S1) and (S2) of Theorem 3.1 are satisfied. Now, we check that (S3) also holds. Suppose that $\|\varkappa(t)\| \leq \varepsilon$, then,

$$\begin{aligned} |\varkappa(t)| &= \left| \frac{e^{-2t}}{3} + \frac{\ln(1 + |\varkappa(\sqrt{t})|)}{4 + t^2} + \frac{|\varkappa(\sqrt{t})|}{1 + |\varkappa(\sqrt{t})|} \right. \\ &\quad \left. + \frac{1}{\Gamma_{1/4}(\frac{5}{2})} \int_0^{t^2} \left(t^2 - \frac{1}{4}s\right)^{3/2} \sqrt{\frac{1 + \varkappa(\sqrt{t})}{1 + st}} d_qs \right| \\ &\leq \frac{1}{3} + \frac{1}{4}\varepsilon + \frac{1}{3} + \frac{1}{\Gamma_{1/4}(\frac{5}{2})} B_{1/4} \left(\frac{5}{2}, 1\right) \sqrt{1 + \varepsilon} \leq \varepsilon. \end{aligned} \quad (4.6)$$

Hence, (S3) holds if, $\varepsilon \geq 2.67961$. This implies that the equation has at least one solution in $C(I_1)$.

Remark 2. Since there is no constants c_1 and c_2 satisfying the inequalities (Sublinear condition), $|p(t, s, x)| \leq c_1 + c_2|x|$, for all $t, s \in I_a$ and $x \in \mathbb{R}$, the results in [17] and [3] are inapplicable to the q - \mathbb{IE} 4.5.

5 Conclusion

Because many problems of q - \mathbb{IE} s are unsolvable, it is important to ensure that a solution exists in these equations. Therefore, many researchers have published papers in this field and have explained their methods with the results obtained. Accordingly in this paper, the authors presented a new method based on the technique of M.N.C and Petryshyn's F.P.T. The advantages of the proposed method compared to other similar methods are that it has fewer conditions, and there is no need to verify the involved operator maps a closed convex subset onto itself.

References

- [1] N. Adjimi, A. Boutiara, M. E. Samei, S. Etemad and S.Rezapour, On solutions of a hybrid generalized Caputo-type problem via the measure of non-compactness in the generalized version of Darbo's theorem, *J. Ineq. Appl.*, **2023** (2023), 34.
- [2] B. Ahmad, J. J. Nieto, A. Alsaedi and H. Al-Hutami, Existence of solutions for nonlinear fractional q -difference integral equations with two fractional orders and nonlocal four point boundary conditions, *J. Franklin Instit. Engin. Appl. Math.*, **351** (2014), 2890–2909.
- [3] M. Al-Yami, Existence of solutions of q -perturbed quadratic integral equations, *Amer. J. Comput. Math.*, **6** (2016), 166–176.
- [4] A. Alsaedi, B. Ahmad and H. Al-Hutami, A study of nonlinear fractional q -difference equations with nonlocal integral boundary conditions, *Abst. Appl. Anal.*, **8** (2013), 410505, 8 pages.
- [5] J. Alzabut, B. Mohammadaliev and M. E. Samei, Solutions of two fractional q -integro-differential equations under sum and integral boundary value conditions on a time scale, *Adv. Differ. Equat.*, **2020** (2020), 304.
- [6] J. Banas and M. Lecko, Fixed points of the product of operators in Banach algebra, *Panamer. Math. J.*, **12** (2002), 101–109.
- [7] Bhupeshwar, D. Kumar Patel, M. Gabeleh and M. E. Samei, Existence of solutions for Higher order Ψ -Hilfer boundary value problems, *Comput. Appl. Math.*, **43** (2024), 155.
- [8] J. Caballero, A. B. Mingarelli and K. Sadarangani, Existence of solutions of an integral equation of Chandrasekhar type in the theory of radiative transfer, *Elect. J. Diff. Eq.*, **57** (2006), 1–11.
- [9] A. Darbo, Punti uniti in trasformazioni a codominio non compatto, *Rend. Accad. Naz. Lincei.*, **48** (1970), 195–198.
- [10] A. Das, B. Hazarika, V. Parvaneh and M. Mursaleen, Solvability of generalized fractional order integral equations via measures of noncompactness, *Math. Sciences*, **15** (2021), 241–251.
- [11] A. Deep, S. Abbas, B. Sing, M. R. Alharthi and K. S. Nisar, Solvability of functional stochastic integral equations via Darbo's fixed point theorem, *Alex. Engin. J.*, **60**(6) (2021), 5631–5636.
- [12] R. J. Finkelstein, q -gauge theory, *Int. J. Mod. Phys. A*, **11**(4) (1996), 733–746.
- [13] R. J. Finkelstein, Symmetry group of the hydrogen atom, *J. Math. Phys.*, **8**(3) (1967), 443–449.
- [14] L. S. Goldenstein and A. S. Markus, On the measure of non-compactness of bounded sets and of linear operators, *Stud. Alg. Math. Anal.*, Izdat. "Moldovenjaske", Kishinev (1965), 45–54.
- [15] K. N. Ilinski, G. V. Kalinin and A. S. Stepanenko, q -functional field theory for particles with exotic statistics, *Phys. Lett. A*, **232**(6) (1997), 399–408.

-
- [16] F. H. Jackson, q -Difference Equations, The Amer. J. Math., **32**(4) (1910), 305–314.
 - [17] M. Jleli, M. Mursaleen and B. Samet, Q -integral equations of fractional orders, Elect. J. Differ. Equat., **2016**(17) (2016), 1–14.
 - [18] V. Kac and P. Cheung, Quantum Calculus, 2^{nd} edition, Springer, New York, NY, USA, 2002.
 - [19] M. Kazemi and R. Ezzati, Existence of solution for some nonlinear two-dimensional Volterra integral equations via measures of noncompactness, Appl. Math. Comput., **275** (2016), 165–171.
 - [20] M. Kazemi, R. Ezzati and A. Deep, On the solvability of non-linear fractional integral equations of product type, J. Pseudo-Differ. Oper. Appl., **14** (2023), 39.
 - [21] S. Kumar, H. Kumar Singh, B. Singh and V. Arora, Application of petryshyn’s fixed point theorem of existence result for nonlinear 2D volterra functional integral equations, Differ. Equat. Appl., **14**(3) (2022), 487–497.
 - [22] K. Kuratowski, Sur les espaces completes, Fund. Math., **15** (1934), 301–335.
 - [23] X. Li, Z. Han and S. Sun, Existence of positive solutions of nonlinear fractional q -difference equation with parameter, Adv. Differ. Equat., **2013** (2013), 260.
 - [24] M. Metwali and K. Cicho’n, On solutions of some delay Volterra integral problems on a half line, Nonlinear Anal. Model., **26** (2021), 661–677.
 - [25] L. N. Mishra, R. P. Agarwal and M. Sen, Solvability and asymptotic behavior for some nonlinear quadratic integral equation involving Erdelyi-Kober fractional integrals on the unbounded interval, Prog. Fract. Differ. and Appl., **2** (2016), 153–168.
 - [26] H. Nashine, R. Arab and R. Agarwal, Existence of solutions of system of functional integral equations using measure of noncompactness, Int. J. Nonlinear Anal., **12** (2021), 583–595.
 - [27] W. V. Petryshyn, Structure of the fixed points sets of k -set contractions, Arch. Rational Mech. Anal., **40** (1971), 312–328.
 - [28] N. Pongarm, S. Asawasamrit, J. Tariboon and S. K. Ntouyas, Multi strip fractional q -integral boundary value problems for nonlinear fractional q -difference equations, Adv. Differ. Equat., **2014** (2014), 1–17.
 - [29] P. Ramesh Patle, M. Gabeleh, V. Rakočević and M. E. Samei, New best proximity point (pair) theorems via MNC and application to the existence of optimum solutions for a system of ψ -Hilfer fractional differential equations, Revi. Real Acad. Cienc. Exact., Físi. Natural. Serie A. Matem., **117** (2023), 124.
 - [30] M. E. Samei, Existence of solutions for a system of singular sum fractional q -differential equations via quantum calculus, Adv. Differ. Equat., **2020** (2020), 23.
 - [31] M. E. Samei, Employing Kuratowski measure of non-compactness for positive solutions of system of singular fractional q -differential equations with numerical effects, Filomat, **34**(9) (2020), 1–19.

- [32] M. E. Samei and W. Yang, Existence of solutions for k -dimensional system of multi-term fractional q -integro-differential equations under anti-periodic boundary conditions via quantum calculus, *Math. Meth. Appl. Sci.*, **43**(7) (2020), 4360–4382.
- [33] M. Sen, D. Saha and R. P. Agarwal, A Darbo fixed point theory approach towards the existence of a functional integral equation in a Banach algebra, *Appl. Math. Comput.*, **358** (2019), 111–118.
- [34] H. M. Srivastava, A. Das, B. Hazarika and S. A. Mohiuddine, Existence of solutions for nonlinear functional integral equation of two variables in Banach Algebra, *Symmetry*, **11** (2019), 674.

Hamid Reza Sahebi Department of Mathematics, Ashtian Branch, Islamic Azad University, Ashtian, Iran

E-mail: sahebi.aiau.ac.ir@gmail.com

Manochehr Kazemi Department of Mathematics, Ashtian Branch, Islamic Azad University, Ashtian, Iran, and Department of Mathematics, Central Tehran Branch, Islamic Azad University, tehran, Iran,

E-mail: univer_ka@yahoo.com