



Inverse problem for Dirac operators with a small delay

Nebojša Djurić and Biljana Vojvodić

Abstract. This paper addresses inverse spectral problems associated with Dirac-type operators with a constant delay, specifically when this delay is less than one-third of the interval length. Our research focuses on eigenvalue behavior and operator recovery from spectra. We find that two spectra alone are insufficient to fully recover the potentials. Additionally, we consider the Ambarzumian-type inverse problem for Dirac-type operators with a delay. Our results have significant implications for the study of inverse problems related to differential operators with constant delay and may inform future research directions in this field.

Keywords. Dirac-type operator, constant delay, functional-differential operator, inverse spectral problem

1 Introduction

The main goal of this paper is to study a two-spectra inverse problem of recovering a system of differential equations with delay. Specifically, we consider the boundary value problems $\mathcal{B}_j(a, p, q)$, $j = 1, 2$, for Dirac-type system of the form

$$\begin{aligned} By'(x) + Q(x)y(x-a) &= \lambda y(x), \quad 0 < x < \pi, \\ y_1(0) = y_2(\pi) &= 0, \end{aligned} \tag{1.1}$$

where

$$B = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \quad y(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \end{bmatrix}, \quad Q(x) = \begin{bmatrix} p(x) & q(x) \\ q(x) & -p(x) \end{bmatrix},$$

as $p(x), q(x) \in L_2(0, \pi)$ are complex-valued functions with $Q(x) = 0$ on $(0, a)$ and the delay $a \in (0, \frac{\pi}{3})$. Let $\{\lambda_{n,j}\}_{n \in \mathbb{Z}}$, $j = 1, 2$, be the eigenvalues of $\mathcal{B}_j(a, p, q)$. The inverse problem is formulated as follows.

Inverse problem 1. Given the two spectra $\{\lambda_{n,j}\}_{n \in \mathbb{Z}}$, $j = 1, 2$, construct $p(x)$, $q(x)$.

Received date: August 2, 2024; Published online: March 12, 2025.
 2010 *Mathematics Subject Classification.* 34A55, 34K29.
 Corresponding author: Nebojša Djurić.

The idea of using two sequences of eigenvalues for a Dirac-type system on a finite interval originated with Gasymov's and Dzabiev's classic paper [14], where they developed a constructive solution based on transformation operators and obtained the characterization of the spectral data. Two-spectra inverse problems for Dirac-type systems have been considered in [1, 11]. Research on inverse problems for the Dirac operator with a delay started a few years ago, and the first result in this direction is the paper [4]. In that paper, the authors restrict themselves to the case of a "large delay" when the dependence of the characteristic functions on the potential is linear. However, for the considered case, they achieved their objectives by answering a full range of questions regarding uniqueness, solvability, and uniform stability. Very recently, progress has been made in the nonlinear case in the paper [10], by proving that two spectra uniquely determine potentials if $a \in [2\pi/5, \pi/2)$, yet it is not possible in the case when $a \in [\pi/3, 2\pi/5)$. In [26], more research has been conducted for the case where $a \in [2\pi/5, \pi/2)$ by giving a necessary and sufficient condition for the solvability and formulating the stability of the inverse problem. The question of whether two spectra are enough to uniquely recover potentials for $a < \pi/3$ remained unanswered. In this paper, we fill this gap by giving a negative answer to the question of whether the unique construction of the potentials is possible from the two spectra.

The answer to this question will also be valuable in addressing similar issues with other types of differential operators with delays. A major avenue for further research involves boundary value problems associated with differential equations that include two delays. In the papers [21], [25] authors study four boundary value problems $\mathcal{D}_j(P, Q, m)$, $m = 0, 1$, $j = 1, 2$, for Dirac-type system of the form

$$By'(x) + (-1)^m P(x)y(x - a_1) + Q(x)y(x - a_2) = \lambda y(x), \quad 0 < x < \pi, \quad (1.2)$$

$$y_1(0) = y_j(\pi) = 0,$$

where $\frac{\pi}{3} \leq a_1 < a_2 < \pi$,

$$P(x) = \begin{bmatrix} p_1(x) & p_2(x) \\ p_2(x) & -p_1(x) \end{bmatrix}, \quad Q(x) = \begin{bmatrix} q_1(x) & q_2(x) \\ q_2(x) & -q_1(x) \end{bmatrix},$$

and $p_1(x), p_2(x), q_1(x), q_2(x) \in L^2[0, \pi]$ are complex-valued functions such that

$$P(x) = 0, \quad x \in (0, a_1), \quad Q(x) = 0, \quad x \in (0, a_2).$$

Let $\{\lambda_{n,j}^m\}$ be the spectrum of the boundary value problem $\mathcal{D}_j(P, Q, m)$, $j = 1, 2$, $m = 0, 1$, and assume that delays a_1 and a_2 are known. In the papers [21] and [25] the authors consider the following inverse problem.

Inverse problem 2. Given the four spectra $\{\lambda_{n,j}^m\}_{n \in \mathbb{Z}}$, $j = 1, 2$, $m = 0, 1$, construct the matrix functions $P(x)$ and $Q(x)$.

It is obvious that equation (1.2) becomes equation (1.1) when $P(x) = 0$, and then Inverse problem 2 reduces to Inverse problem 1. In the paper [25] it has been shown that Inverse problem 2 has a unique solution as soon as the delays $a_1, a_2 \in [2\pi/5, \pi)$. One can easily conclude that the solution of Inverse Problem 2 is not unique in the case $a_2 \in [\pi/3, 2\pi/5)$. In fact, it has been shown in the paper [10] that the solution of Inverse problem 1 is not unique for $a \in [\pi/3, 2\pi/5)$. Therefore, taking $P(x) = 0$ in equation (1.2), we obtain the same conclusion for Inverse Problem 2. The case $a_1 \in [\pi/3, 2\pi/5)$ and $a_2 \in [2\pi/5, \pi)$ has been considered in the paper [21]. It has been established that Inverse problem 2 possesses a unique solution in the case where $2a_1 + a_2/2 \geq \pi$, whereas no uniqueness holds if $2a_1 + a_2/2 < \pi$.

In the paper [27], the authors introduced Dirac-type operators with a global constant delay on a star graph consisting of m equal edges. They proved that the uniqueness theorem is valid in the case when the delay is greater than one-half of the length of the interval.

Many of the differential equations with delays, primarily sourced from the literature in biological sciences, await exploration. These encompass different types of models spanning population biology, physiology, epidemiology, economics, neural networks, and intricate control mechanisms of mechanical systems. The application of differential equations with delay is covered in the books by Erneux [12], Polyanin [18], and Smith [19]. Additionally, Norkin [15] dealt with models involving the Sturm-Liouville equation with delay,

$$-y''(x) + q(x)y(x-a) = \lambda y(x), \quad 0 < x < \pi.$$

For this reason, the inverse spectral problems for Sturm-Liouville operators with delay have been studied in detail. Some important results in this field can be found in [2, 5, 7, 8, 9, 13, 16, 17, 20]. In addition, inverse problems for Sturm-Liouville operators with two delays have been thoroughly investigated in articles [22, 23, 24].

The paper is organized as follows. In Section 2, we describe characteristic functions and study the asymptotic behavior of eigenvalues. In Section 3, we construct the class of iso-bispectral potentials $p(x), q(x)$ that have the same characteristic functions. In this way, we show that two spectra of boundary value problems $\mathcal{B}_j(a, p, q)$, $j = 1, 2$, are not enough to uniquely determine the potentials $p(x), q(x)$. In Section 4, we study the Ambarzumian-type inverse problem for Dirac operators with constant delay.

2 Characteristic functions and spectra

Let $a \in [\pi/(N+1), \pi/N]$ for $N \in \mathbb{N}$ and $S(x, \lambda) = \begin{bmatrix} s_1(x, \lambda) \\ s_2(x, \lambda) \end{bmatrix}$ be the fundamental (vector) solution of Eq. (1.1) satisfying the initial condition at the origin:

$$S(0, \lambda) = \begin{bmatrix} 0 \\ -1 \end{bmatrix}.$$

The solution $S(x, \lambda)$ is the unique solution of the following integral equation

$$S(x, \lambda) = S_0(x, \lambda) + \int_a^x Q(t)B^{-1}(\lambda(x-t))S(t-a, \lambda) dt \quad (2.1)$$

where

$$S_0(x, \lambda) = \begin{bmatrix} \sin \lambda x \\ -\cos \lambda x \end{bmatrix}, \quad B(t) = \begin{bmatrix} \sin t & \cos t \\ -\cos t & \sin t \end{bmatrix}.$$

The method of successive approximations gives

$$\begin{aligned} S(x, \lambda) &= \sum_{k=0}^N S_k(x, \lambda), \\ S_k(x, \lambda) &= \int_{ka}^x Q(t)B^{-1}(\lambda(x-t))S_{k-1}(t-a, \lambda) dt, \quad x > ka, \\ S_k(x, \lambda) &= 0, \quad x \leq ka, \quad k = \overline{1, N}. \end{aligned} \quad (2.2)$$

According to (2.2), we can calculate (see [10])

$$S_1(x, \lambda) = \int_a^x Q(t_1) \begin{bmatrix} \cos \lambda(x - 2t_1 + a) \\ -\sin \lambda(x - 2t_1 + a) \end{bmatrix} dt_1,$$

$$S_2(x, \lambda) = \int_{2a}^x Q(t_1) \int_a^{t_1-a} Q(t_2) \begin{bmatrix} \sin \lambda(x - 2t_1 + 2t_2) \\ -\cos \lambda(x - 2t_1 + 2t_2) \end{bmatrix} dt_2 dt_1.$$

In a similar way, we obtain

$$S_3(x, \lambda) = \int_{3a}^x Q(t_1) \int_{2a}^{t_1-a} Q(t_2) \int_a^{t_2-a} Q(t_3) \begin{bmatrix} \cos \lambda(x - 2t_1 + 2t_2 - 2t_3 + a) \\ -\sin \lambda(x - 2t_1 + 2t_2 - 2t_3 + a) \end{bmatrix} dt_3 dt_2 dt_1,$$

and consequently, by induction, we determine the form of the functions $S_k(x, \lambda)$ for the case $k = n$, $n \in \mathbb{N}$ is even, as

$$S_n(x, \lambda) = \int_{na}^x Q(t_1) dt_1 \int_{(n-1)a}^{t_1-a} Q(t_2) dt_2 \cdots \int_a^{t_{n-1}-a} Q(t_n) \begin{bmatrix} \sin \lambda(x - 2t_1 + 2t_2 - 2t_3 + \dots - 2t_{n-1} + 2t_n) \\ -\cos \lambda(x - 2t_1 + 2t_2 - 2t_3 + \dots - 2t_{n-1} + 2t_n) \end{bmatrix} dt_n, \quad (2.3)$$

and for the case $k = m$, $m \in \mathbb{N}$ is odd, as

$$S_m(x, \lambda) = \int_{ma}^x Q(t_1) dt_1 \int_{(m-1)a}^{t_1-a} Q(t_2) dt_2 \cdots \int_a^{t_{m-1}-a} Q(t_m) \begin{bmatrix} \cos \lambda(x - 2t_1 + 2t_2 - 2t_3 + \dots + 2t_{m-1} - 2t_m + a) \\ -\sin \lambda(x - 2t_1 + 2t_2 - 2t_3 + \dots + 2t_{m-1} - 2t_m + a) \end{bmatrix} dt_m. \quad (2.4)$$

The functions

$$\Delta_j(\lambda) := s_j(\pi, \lambda), \quad j = 1, 2, \quad (2.5)$$

are called the *characteristic functions* of the boundary value problems $\mathcal{B}_j(a, p, q)$. These functions are entire in λ and their zeros coincide with the eigenvalues of $\mathcal{B}_j(a, p, q)$.

In order to express the characteristic functions in a more convenient way, let us introduce the notation for the functions $S_k(x, \lambda)$ given in Eq. (2.2):

$$S_k(x, \lambda) = \begin{bmatrix} s_{1,k}(x, \lambda) \\ s_{2,k}(x, \lambda) \end{bmatrix}, \quad k = \overline{1, N}.$$

Taking Eq.s (2.1), (2.3), (2.4), and (2.5) into account, we can rewrite the characteristic functions as

$$\Delta_j(\lambda) = s_j(\pi, \lambda) = s_{j,0}(\pi, \lambda) + \sum_{k=1}^N s_{j,k}(\pi, \lambda), \quad j = 1, 2. \quad (2.6)$$

Now, let $k = \overline{1, N}$. Then, for the case $k = m$, m is odd, the characteristic functions become

$$\begin{aligned} s_{j,m}(\pi, \lambda) = & \int_{ma}^{\pi} dt_1 \int_{(m-1)a}^{t_1-a} dt_2 \cdots \int_a^{t_{m-1}-a} \left(Q_{j,1}^m(t_1, t_2, \dots, t_m) \right. \\ & \left. \cos \lambda \left(\pi - 2t_1 + 2t_2 - 2t_3 + \cdots + 2t_{m-1} - 2t_m + a \right) \right) dt_m \\ & - \int_{ma}^{\pi} dt_1 \int_{(m-1)a}^{t_1-a} dt_2 \cdots \int_a^{t_{m-1}-a} Q_{j,2}^m(t_1, t_2, \dots, t_m) \\ & \left. \sin \lambda \left(\pi - 2t_1 + 2t_2 - 2t_3 + \cdots + 2t_{m-1} - 2t_m + a \right) \right) dt_m, \end{aligned} \quad (2.7)$$

and for the case $k = n$, n is even,

$$\begin{aligned} s_{j,n}(\pi, \lambda) = & \int_{na}^{\pi} dt_1 \int_{(n-1)a}^{t_1-a} dt_2 \cdots \int_a^{t_{n-1}-a} \left(Q_{j,1}^n(t_1, t_2, \dots, t_n) \right. \\ & \left. \sin \lambda \left(\pi - 2t_1 + 2t_2 - 2t_3 + \cdots - 2t_{n-1} + 2t_n \right) \right) dt_n \\ & - \int_{na}^{\pi} dt_1 \int_{(n-1)a}^{t_1-a} dt_2 \cdots \int_a^{t_{n-1}-a} Q_{j,2}^n(t_1, t_2, \dots, t_n) \\ & \left. \cos \lambda \left(\pi - 2t_1 + 2t_2 - 2t_3 + \cdots - 2t_{n-1} + 2t_n \right) \right) dt_n. \end{aligned} \quad (2.8)$$

where we use the notation $Q_{j,l}^k(t_1, t_2, \dots, t_k)$ for the entries in the j -th row and l -th column of the matrix

$$Q^k(t_1, t_2, \dots, t_k) := Q(t_1)Q(t_2) \cdots Q(t_k).$$

One can easily show that next relations hold:

$$\begin{aligned} Q_{11}^k(t_1, t_2, \dots, t_k) &= (-1)^k Q_{22}^k(t_1, t_2, \dots, t_k), \\ Q_{21}^k(t_1, t_2, \dots, t_k) &= (-1)^{k+1} Q_{12}^k(t_1, t_2, \dots, t_k). \end{aligned} \quad (2.9)$$

It is obvious from (2.7) and (2.8) that the next asymptotic holds

$$s_{j,k}(\pi, \lambda) = O\left(\exp(|\operatorname{Im} \lambda|(\pi - ka))\right), \quad |\lambda| \rightarrow \infty. \quad (2.10)$$

Taking equations (2.6), (2.7), (2.8) and (2.10) into account, we derive the asymptotics of the characteristic functions given in Eq. (2.6), for $|\lambda| \rightarrow \infty$,

$$\begin{aligned} \Delta_1(\lambda) = & \sin \lambda \pi + \int_a^{\pi} p(t) \cos \lambda(\pi - 2t_1 + a) dt_1 \\ & - \int_a^{\pi} q(t) \sin \lambda(\pi - 2t_1 + a) dt_1 + O\left(\exp(|\operatorname{Im} \lambda|(\pi - 2a))\right), \end{aligned} \quad (2.11)$$

$$\begin{aligned} \Delta_2(\lambda) = & -\cos \lambda \pi + \int_a^{\pi} q(t) \cos \lambda(\pi - 2t_1 + a) dt_1 \\ & + \int_a^{\pi} p(t) \sin \lambda(\pi - 2t_1 + a) dt_1 + O\left(\exp(|\operatorname{Im} \lambda|(\pi - 2a))\right). \end{aligned} \quad (2.12)$$

Therefore, the following theorem can be proved using the standard method [5, Lemma 3], Rouché's theorem, and Eqs (2.11), (2.12).

Theorem 2.1. *The boundary value problems $\mathcal{B}_j(a, p, q)$, $j = 1, 2$, have infinitely many eigenvalues $\{\lambda_{n,j}\}_{n \in \mathbb{Z}}$ of the form*

$$\lambda_{n,j} = n + \frac{1-j}{2} + o(1), \quad |n| \rightarrow \infty.$$

Lemma 2.1. *The specification of the spectrum $\{\lambda_{n,j}\}_{n \in \mathbb{Z}}$, $j = 1, 2$, uniquely determines the characteristic function $\Delta_j(\lambda)$ by the formulae*

$$\Delta_1(\lambda) = \pi(\lambda_{0,1} - \lambda) \prod_{|n| \in \mathbb{N}} \frac{\lambda_{n,1} - \lambda}{n} \exp\left(\frac{\lambda}{n}\right), \quad \Delta_2(\lambda) = \prod_{n \in \mathbb{Z}} \frac{\lambda_{n,2} - \lambda}{n - 1/2} \exp\left(\frac{\lambda}{n - 1/2}\right).$$

Proof. The proof can be performed similarly as in [3, Theorem 5] using the Hadamard factorization method. \square

In addition to examining the asymptotic behavior of the characteristic functions, solving the Ambarzumian-type inverse problem requires considering appropriate combinations of these functions and determining their asymptotic behavior. For that purpose, let us introduce the functions:

$$\mathcal{L}(\lambda) = \frac{1}{2} \left(\Delta_1(\lambda) + \Delta_1(-\lambda) + i(\Delta_2(\lambda) - \Delta_2(-\lambda)) \right), \quad (2.13)$$

$$\mathcal{M}(\lambda) = \exp(i\lambda\pi) + \frac{1}{2} \left(\Delta_2(\lambda) + \Delta_2(-\lambda) + i(-\Delta_1(\lambda) + \Delta_1(-\lambda)) \right). \quad (2.14)$$

It follows from Eq.s (2.6), (2.7), (2.8), (2.9), (2.13) and (2.14) that

$$\mathcal{L}(\lambda) = \sum_{k=1}^N \mathcal{L}_k(\lambda), \quad \mathcal{M}(\lambda) = \sum_{k=1}^N \mathcal{M}_k(\lambda),$$

holds for

$$\begin{aligned} \mathcal{L}_k(\lambda) = & \int_{ka}^{\pi} dt_1 \int_{(k-1)a}^{t_1-a} dt_2 \cdots \int_a^{t_{k-1}-a} \left(Q_{2-\varphi(k),1}^k(t_1, t_2, \dots, t_k) \right. \\ & \left. \exp i\lambda(\pi - 2t_1 + 2t_2 - 2t_3 + \cdots + (-1)^{k-1}2t_{k-1} + (-1)^k 2t_k + a\varphi(k)) \right) dt_k, \end{aligned} \quad (2.15)$$

$$\begin{aligned} \mathcal{M}_k(\lambda) = & (-1)^{k+1} \int_{ka}^{\pi} dt_1 \int_{(k-1)a}^{t_1-a} dt_2 \cdots \int_a^{t_{k-1}-a} \left(Q_{1+\varphi(k),1}^k(t_1, t_2, \dots, t_k) \right. \\ & \left. \exp i\lambda(\pi - 2t_1 + 2t_2 - 2t_3 + \cdots + (-1)^{k-1}2t_{k-1} + (-1)^k 2t_k + a\varphi(k)) \right) dt_k, \end{aligned} \quad (2.16)$$

where $\varphi(2k) = 0$ and $\varphi(2k-1) = 1$, $k \in \mathbb{N}$. In particular, the following asymptotic formulas are valid for $\text{Im}\lambda > 0$ as $|\lambda| \rightarrow \infty$:

$$\mathcal{L}_k(\lambda) = O(\exp(-i\lambda(\pi - ka))), \quad \mathcal{M}_k(\lambda) = O(\exp(-i\lambda(\pi - ka))).$$

3 Iso-bispectral potentials

In this Section we shall establish non-uniqueness of the solution of Inverse problem 1. Specifically, we construct an infinite family of iso-bispectral potentials $p_\alpha(x)$ and $q_\beta(x)$, $\alpha, \beta \in \mathbb{C}$, i.e. for which both problems $\mathcal{B}_1(a, p_\alpha, q_\beta)$ and $\mathcal{B}_2(a, p_\alpha, q_\beta)$ possess one and the same pair of spectra. For that purpose we use the ideas of the construction of the counterexample from the papers [9] and [10]. We restrict ourselves with potentials vanishing on the interval $(3a, \pi)$ and using (2.6, 2.7, 2.8) we obtain the following representation of the characteristic functions

$$\begin{aligned}\Delta_1(\lambda) &:= \sin \lambda \pi + \int_a^{3a} p(t) \cos \lambda(\pi - 2t + a) dt - \int_a^{3a} q(t) \sin \lambda(\pi - 2t + a) dt \\ &\quad + \int_{2a}^{3a} \int_a^{t-a} (p(t)p(s) + q(t)q(s)) \sin \lambda(\pi - 2t + 2s) ds dt \\ &\quad + \int_{2a}^{3a} \int_a^{t-a} (q(t)p(s) - p(t)q(s)) \cos \lambda(\pi - 2t + 2s) ds dt, \\ \Delta_2(\lambda) &:= -\cos \lambda \pi + \int_a^{3a} p(t) \sin \lambda(\pi - 2t + a) dt + \int_a^{3a} q(t) \cos \lambda(\pi - 2t + a) dt \\ &\quad - \int_{2a}^{3a} \int_a^{t-a} (p(t)p(s) + q(t)q(s)) \cos \lambda(\pi - 2t + 2s) ds dt, \\ &\quad + \int_{2a}^{3a} \int_a^{t-a} (q(t)p(s) - p(t)q(s)) \sin \lambda(\pi - 2t + 2s) ds dt.\end{aligned}$$

Let us assume that the delay a is known. For fixed $a \in (0, \frac{\pi}{3})$ let us define the integral operator

$$M_h f(x) = \int_{3a/2}^{7a/2-x} f(t) h\left(t + x - \frac{a}{2}\right) dt, \quad x \in \left(\frac{3a}{2}, 2a\right),$$

for some non-zero real function $h(x) \in L_2(\frac{5a}{2}, 3a)$.

Operator M_h is a non-zero compact Hermitian operator in $L_2(\frac{3a}{2}, 2a)$ and hence, it has at least one non-zero eigenvalue η . Putting $h_m(x) = (-1)^m h(x)/\eta$ for $m \in \{0, 1\}$, we obtain that $(-1)^m$ is an eigenvalue of the operator M_{h_m} . Let $e_m(x)$ be a corresponding eigenfunction, i.e.

$$M_{h_m}(x) = (-1)^m e_m(x), \quad x \in \left(\frac{3a}{2}, 2a\right).$$

Now we construct the iso-bispectral family of functions

$$D = \{p_\alpha(x), q_\beta(x) : \alpha, \beta \in \mathbb{C}\}$$

where

$$p_\alpha(x) = \begin{cases} 0, & x \in (0, \frac{3a}{2}) \cup (2a, \pi), \\ \alpha e_1(x), & x \in (\frac{3a}{2}, 2a), \end{cases} \quad (3.1)$$

and

$$q_\beta(x) = \begin{cases} 0, & x \in (0, \frac{3a}{2}) \cup (2a, \frac{5a}{2}) \cup (3a, \pi), \\ \beta e_0(x), & x \in (\frac{3a}{2}, 2a), \\ h(x), & x \in (\frac{5a}{2}, 3a). \end{cases} \quad (3.2)$$

We shall prove that the spectra of the boundary value problem $\mathcal{B}_j(a, p_\alpha, q_\beta)$ for $j = 1, 2$, is independent of α and β , hence, the solution of the Inverse problem 1 is not unique for a delay less than one-third of the interval length.

Theorem 3.1. *Let delay $a \in (0, \frac{\pi}{3})$. The spectra $\{\lambda_{n,j}\}_{n \in \mathbb{Z}}$ of the boundary value problem $\mathcal{B}_j(a, p_\alpha, q_\beta)$ for $j = 1, 2$, are independent of α and β .*

Proof. Taking into account that all multiple integrals except the double one in (2.7) and (2.8) are equal to zero, we obtain that the characteristic functions for the boundary value problem $\mathcal{B}_j(a, p_\alpha, q_\beta)$ have the form

$$\Delta_1(\lambda) = \sin \lambda \pi + \int_{a/2}^{5a/2} K_1(x) \cos \lambda(\pi - 2x) dx - \int_{a/2}^{5a/2} K_2(x) \sin \lambda(\pi - 2x) dx, \quad (3.3)$$

$$\Delta_2(\lambda) = -\cos \lambda \pi + \int_{a/2}^{5a/2} K_1(x) \sin \lambda(\pi - 2x) dx + \int_{a/2}^{5a/2} K_2(x) \cos \lambda(\pi - 2x) dx, \quad (3.4)$$

where

$$K_1(x) = \begin{cases} p(x + a/2), & x \in (a/2, a) \cup (2a, 5a/2), \\ p(x + a/2) - \int_{x+a}^{3a} p(t)q(t-x)dt + \int_{x+a}^{3a} q(t)p(t-x)dt, & x \in (a, 2a), \end{cases}$$

$$K_2(x) = \begin{cases} q(x + a/2), & x \in (a/2, a) \cup (2a, 5a/2), \\ q(x + a/2) - \int_{x+a}^{3a} p(t)p(t-x)dt - \int_{x+a}^{3a} q(t)q(t-x)dt, & x \in (a, 2a). \end{cases}$$

Then, from (3.1) and (3.2) we obtain

$$K_1(x - a/2) = \begin{cases} 0, & x \in (0, \frac{3a}{2}) \cup (2a, 3a), \\ p_\alpha(x) + \int_{3a/2}^{7a/2-x} p_\alpha(t)h(t+x-a/2)dt, & x \in (\frac{3a}{2}, 2a), \end{cases}$$

$$K_2(x - a/2) = \begin{cases} 0, & x \in (0, \frac{3a}{2}) \cup (2a, \frac{5a}{2}), \\ q_\beta(x) - \int_{3a/2}^{7a/2-x} q_\beta(t)h(t+x-a/2)dt, & x \in (\frac{3a}{2}, 2a), \\ h(x), & x \in (\frac{5a}{2}, 3a). \end{cases}$$

Since $e_0(x)$ and $e_1(x)$ are eigenfunctions of the operator M_h corresponding to the eigenvalues 1 and -1 respectively, we obtain

$$K_1(x - a/2) = 0, \quad x \in (a, 3a),$$

and

$$K_2(x - a/2) = \begin{cases} 0, & x \in (a, \frac{5a}{2}), \\ h(x), & x \in (\frac{5a}{2}, 3a). \end{cases}$$

Then, using (3.3) and (3.4) we conclude that characteristic functions for family of iso-bispectral potentials D have the form

$$\Delta_1(\lambda) = \sin \lambda \pi - \int_{5a/2}^{3a} h(x) \sin \lambda(\pi - 2x + a) dx,$$

$$\Delta_2(\lambda) = -\cos \lambda \pi + \int_{5a/2}^{3a} h(x) \cos \lambda(\pi - 2x + a) dx,$$

i.e. they are independent of α and β . \square

4 Ambarzumian-type inverse problem

In Section 3, we proved that the solution to Inverse Problem 1 is not unique. Using the results from Section 3 and the paper [10], we conclude that two spectra are not sufficient to recover the potentials when $a \in (0, 2\pi/5)$. However, it is interesting to examine whether the uniqueness of the solution holds for the Ambarzumian-type inverse problem for Dirac operators with delay. In the papers [5, 13], the authors prove that the uniqueness of the solution holds for the Ambarzumian-type inverse problem for the Sturm-Liouville operators with delay. For this reason, we consider the boundary value problems $\tilde{\mathcal{B}}_j = \mathcal{B}_j(\tilde{p}, \tilde{q}, a)$, where $\tilde{p}(x) = \tilde{q}(x) = 0$ for $j = 1, 2$. Since it is known that for $a \in [2\pi/5, \pi)$, two spectra uniquely determine the potentials $p(x)$ and $q(x)$ (see [4, 10]), we will further consider the case $a \in (0, 2\pi/5)$.

Let $\{\tilde{\lambda}_{n,j}\}_{n \in \mathbb{Z}}$ be the eigenvalues of the boundary value problem $\tilde{\mathcal{B}}_j = \mathcal{B}_j(\tilde{p}, \tilde{q}, a)$, $j = 1, 2$. Denote by $\tilde{\Delta}_j(\lambda)$ the characteristic function of $\tilde{\mathcal{B}}_j$, $j = 1, 2$. From (2.6), (2.7) and (2.8) we obtain $\tilde{\Delta}_1(\lambda) = \sin \lambda \pi$ and $\tilde{\Delta}_2(\lambda) = -\cos \lambda \pi$. Therefore, it is obvious that $\tilde{\lambda}_{n,j} = n - j/2$, $n \in \mathbb{Z}$, $j = 1, 2$.

Theorem 4.1. *Let the delay $a \in (0, 2\pi/5)$ be known. If $\lambda_{n,j} = \tilde{\lambda}_{n,j}$ for all $n \in \mathbb{Z}$, $j = 1, 2$, then $p(x) = q(x) = 0$ a.e. on (a, π) .*

Proof. Let us assume that $\lambda_{n,j} = \tilde{\lambda}_{n,j}$, $n \in \mathbb{Z}$, $j = 1, 2$, holds. By virtue of Lemma 2.1 one has

$$\Delta_1(\lambda) = \sin \lambda \pi, \quad \Delta_2(\lambda) = -\cos \lambda \pi,$$

and consequently $\mathcal{L}(\lambda) = 0$ and $\mathcal{M}(\lambda) = 0$. From (2.13) it follows that

$$\mathcal{L}_1(\lambda) = -\mathcal{L}^+(\lambda), \quad \mathcal{M}_1(\lambda) = -\mathcal{M}^+(\lambda), \quad (4.1)$$

where for $k \geq 2$,

$$\mathcal{L}^+(\lambda) = \sum_{k=2}^N \mathcal{L}_k(\lambda), \quad \mathcal{M}^+(\lambda) = \sum_{k=2}^N \mathcal{M}_k(\lambda),$$

and for $k = 1$, $\mathcal{L}^+(\lambda) = 0$, $\mathcal{M}^+(\lambda) = 0$. Let notice that

$$\begin{aligned} \mathcal{L}_1(\lambda) &= \int_a^\pi p(t_1) \exp(i\lambda(\pi - 2t_1 + a)) dt_1, \\ \mathcal{M}_1(\lambda) &= \int_a^\pi q(t_1) \exp(i\lambda(\pi - 2t_1 + a)) dt_1. \end{aligned} \quad (4.2)$$

Now we divide the proof of the theorem in three steps.

Step (1): Consider the case when $p(x) = q(x) = 0$ a.e. on the interval $(2a, \pi)$. In this scenario, we have $\mathcal{L}^+(\lambda) = 0$ and $\mathcal{M}^+(\lambda) = 0$. From equation (4.1), it follows that $\mathcal{L}_1(\lambda) = \mathcal{M}_1(\lambda) = 0$. Consequently, from equation (4.2), we deduce that $p(x) = q(x) = 0$ a.e. on the interval (a, π) .

Step (2): In this step, we observe the case when $p(x) = q(x) = 0$ a.e. on the interval $(\pi - \nu a/2, \pi)$ for a fixed $\nu = 0, 2N - 3$. We will show that in this case $p(x) = q(x) = 0$ a.e. on the interval $(\pi - (\nu + 1)a/2, \pi)$. We consider the case $\pi - \nu a/2 > 2a$; otherwise, we arrive at Step (1) and the proof is complete. Then from (2.15, 2.16) we obtain

$$\begin{aligned}\mathcal{L}_2(\lambda) &= \int_{2a}^{\pi - \nu a/2} dt_1 \int_a^{t_1 - a} Q_{2,1}^2(t_1, t_2) \exp(-i\lambda(-\pi + 2t_1 - 2t_2)) dt_2, \\ \mathcal{M}_2(\lambda) &= \int_{2a}^{\pi - \nu a/2} dt_1 \int_a^{t_1 - a} Q_{1,1}^2(t_1, t_2) \exp(-i\lambda(-\pi + 2t_1 - 2t_2)) dt_2\end{aligned}$$

In the last integrals we have that $-\pi + 2t_1 - 2t_2 \in (2a - \pi, \pi - (\nu + 2)a)$ and for $\text{Im}\lambda > 0$, $|\lambda| \rightarrow \infty$, the next asymptotic formulas hold:

$$\begin{aligned}\mathcal{L}_2(\lambda) &= O(\exp(-i\lambda(\pi - (\nu + 2)a))), \\ \mathcal{M}_2(\lambda) &= O(\exp(-i\lambda(\pi - (\nu + 2)a))).\end{aligned}\tag{4.3}$$

For $k > 2$, the functions $\mathcal{L}_k(\lambda)$, $\mathcal{M}_k(\lambda)$, have less growth than the right-hand side in (4.3). This means that

$$\begin{aligned}\mathcal{L}^+(\lambda) &= O(\exp(-i\lambda(\pi - (\nu + 2)a))), \\ \mathcal{M}^+(\lambda) &= O(\exp(-i\lambda(\pi - (\nu + 2)a))).\end{aligned}\tag{4.4}$$

It follows from (4.1, 4.2, 4.4) that for $\text{Im}\lambda > 0$, $|\lambda| \rightarrow \infty$,

$$\begin{aligned}\int_a^{\pi - \nu a/2} p(t_1) \exp(-i\lambda(-\pi + 2t_1 - a)) dt_1 &= O(\exp(-i\lambda(\pi - (\nu + 2)a))), \\ \int_a^{\pi - \nu a/2} q(t_1) \exp(-i\lambda(-\pi + 2t_1 - a)) dt_1 &= O(\exp(-i\lambda(\pi - (\nu + 2)a))),\end{aligned}$$

or, which is the same, as

$$\begin{aligned}\exp(i\lambda(2\pi - (\nu + 1)a)) \int_a^{\pi - \nu a/2} p(t_1) \exp(-2i\lambda t_1) dt_1 &= O(1), \\ \exp(i\lambda(2\pi - (\nu + 1)a)) \int_a^{\pi - \nu a/2} q(t_1) \exp(-2i\lambda t_1) dt_1 &= O(1).\end{aligned}\tag{4.5}$$

Let us introduce the functions

$$\begin{aligned}F(\lambda) &= \exp(i\lambda(2\pi - (\nu + 1)a)) \int_{\pi - (\nu + 1)a/2}^{\pi - \nu a/2} p(t_1) \exp(-2i\lambda t_1) dt_1, \\ G(\lambda) &= \exp(i\lambda(2\pi - (\nu + 1)a)) \int_{\pi - (\nu + 1)a/2}^{\pi - \nu a/2} q(t_1) \exp(-2i\lambda t_1) dt_1.\end{aligned}$$

The functions $F(\lambda)$, $G(\lambda)$ are entire in λ . One can show that $F(\lambda) = O(1)$, $G(\lambda) = O(1)$, for $\operatorname{Im} \lambda \leq 0$, $|\lambda| \rightarrow \infty$. On the other hand, from (4.5) follows that $F(\lambda) = O(1)$, $G(\lambda) = O(1)$, for $\operatorname{Im} \lambda > 0$, $|\lambda| \rightarrow \infty$.

By Liouville's theorem [6, page 77], it follows that $F(\lambda) \equiv c_1$ and $G(\lambda) \equiv c_2$, where c_1, c_2 are constants. Since $F(\lambda) = o(1)$, $G(\lambda) = o(1)$ for $\lambda \in \mathbb{R}$ and $|\lambda| \rightarrow \infty$, we conclude that $F(\lambda) \equiv 0$ and $G(\lambda) \equiv 0$. Therefore, it holds

$$\int_{\pi-(\nu+1)a/2}^{\pi-\nu a/2} p(t_1) \exp(-2i\lambda t_1) dt_1 = 0, \quad \int_{\pi-(\nu+1)a/2}^{\pi-\nu a/2} q(t_1) \exp(-2i\lambda t_1) dt_1 = 0.$$

This yields $p(x) = q(x) = 0$ a.e. on interval $(\pi - (\nu + 1)a/2, \pi - \nu a/2)$.

Step (3): Applying Step (2) successively for $\nu = 0, 1, \dots, 2N - 3$, we obtain that $p(x) = q(x) = 0$ a.e. on the interval $(\pi - (N - 1)a, \pi)$. Since $(2a, \pi) \subseteq (\pi - (N - 1)a, \pi)$, from Step (1) we conclude that $p(x) = q(x) = 0$ a.e. on the interval (a, π) . Theorem 4.1 is proved. \square

Acknowledgments

The research is supported by the **Project 19.032/961-68/24**, Ministry of Scientific and Technological Development, Higher Education and Information Society of the Republic of Srpska.

References

- [1] S. Albeverio, R. Hryniv and Ya. V. Mykytyuk, Inverse spectral problems for Dirac operators with summable potentials, *Russian Journal of Mathematical Physics*, **12(4)** (2005), 406–423.
- [2] N. Bondarenko and V. Yurko, An inverse problem for Sturm-Liouville differential operators with deviating argument, *Appl. Math. Lett.*, **83** (2018), 140–144.
- [3] S. Buterin, On the uniform stability of recovering sine-type functions with asymptotically separated zeros, *Math Notes*, **111** (2022), 343–355.
- [4] S. Buterin and N. Djurić, Inverse problems for Dirac operators with constant delay: uniqueness, characterization, uniform stability, *Lobachevskii J. Math.*, **43(6)** (2022), 1492–1501.
- [5] S. A. Buterin, M. Pikula and V. A. Yurko, Sturm-Liouville differential operators with deviating argument, *Tamkang J. Math.*, **48(1)** (2017), 61–71.
- [6] J. B. Conway, *Functions of one complex variable II*, Springer, Graduate texts in mathematics, 1st edition, New York, NY, 1996.
- [7] N. Djurić and S. Buterin, On an open question in recovering Sturm-Liouville-type operators with delay, *Appl. Math. Lett.*, **113** (2021), No 106862.
- [8] N. Djurić and S. Buterin, On non-uniqueness of recovering Sturm-Liouville operators with delay, *Commun. Nonlinear Sci. Numer. Simul.*, **102** (2021), No 105900.

-
- [9] N. Djurić and S. Buterin, Iso-bispectral potentials for Sturm-Liouville-type operators with small delay, *Nonlinear Anal., Real World Appl.*, **63** (2022), No 103390.
 - [10] N. Djurić and B. Vojvodić, Inverse problem for Dirac operators with a constant delay less than half the length of the interval, *Appl. Anal. Discrete Math.*, **17** (2023), 249–261.
 - [11] J. Eckhardt, A. Kostenko and G. Teschl, Inverse uniqueness results for one-dimensional weighted Dirac operators, *Spectral theory and differential equations: V. A. Marchenko's 90th anniversary collection*, 117–133, Providence, RI: American Mathematical Society (AMS), 2014.
 - [12] T. Erneux, *Applied delay differential equations*, Springer, Berlin, 2009.
 - [13] G. Freiling and V. A. Yurko, Inverse problems for Sturm-Liouville differential operators with a constant delay, *Appl. Math. Lett.*, **25**(11) (2012), 1999–2004.
 - [14] M. G. Gasymov and T. T. Džabiev, Solution of the inverse problem by two spectra for the Dirac equation on a finite interval, *Akad. Nauk Azerbaïdžan. SSR Dokl.*, **22**(7) (1966), 3–6.
 - [15] S. B. Norkin, *Differential equations of the second order with retarded argument. Some problems of the theory of vibrations of systems with retardation*. Translated from the Russian by L. J. Grimm and K. Schmitt, *Transl. Math. Monogr.*, **31**, American Mathematical Society (AMS), Providence, RI, 1972.
 - [16] M. Pikula, Determination of a Sturm-Liouville-type differential operator with retarded argument from two spectra, *Mat. Vesnik*, **43**(3-4) (1991), 159–171.
 - [17] M. Pikula, V. Vladičić and B. Vojvodić, Inverse spectral problems for Sturm-Liouville operators with a constant delay less than half the length of the interval and Robin boundary conditions, *Results in Mathematics*, **74**(1) (2019), No 45.
 - [18] A. D. Polyanin, V. G. Sorokin and A. I. Zhurov, *Delay ordinary and partial differential equations*, *Adv. Appl. Math.* (Boca Raton), CRC Press, Boca Raton, FL, 2024.
 - [19] H. L. Smith, *An introduction to delay differential equations with applications to the life sciences*, Springer, New York, **57**, 2011.
 - [20] V. Vladičić, M. Bošković and B. Vojvodić, Inverse problems for Sturm-Liouville-type differential equation with a constant delay under Dirichlet/polynomial boundary conditions, *Bulletin of the Iranian Mathematical Society*, **48**(4) (2022), 1829–1843.
 - [21] B. Vojvodić, N. Djurić and V. Vladičić, On recovering Dirac operators with two delays, *Complex Analysis and Operator Theory*, **18**(5) (2023), No 105.
 - [22] B. Vojvodić, N. Komazec Pavlović and F.A. Çetinkaya, Recovering differential operators with two retarded arguments, *Boletín de la Sociedad Matemática Mexicana*, **28**(3) (2022), No 68.
 - [23] B. Vojvodić and V. Vladičić, Recovering differential operators with two constant delays under Dirichlet/Neumann boundary conditions, *Journal of Inverse and Ill-posed Problems*, **28**(2) (2020), 237–241.
 - [24] B. Vojvodić and V. Vladičić, On recovering Sturm–Liouville operators with two delays, *Journal of Inverse and Ill-posed Problems*, **32**(6) (2024), 1217–1226.

- [25] B. Vojvodić, V. Vladičić and N. Djurić, Inverse problem for Dirac operators with two constant delays, *Journal of Inverse and Ill-posed Problems*, **32(3)** (2024), 573–586.
- [26] F. Wang and C.-F. Yang, Inverse problems for Dirac operators with a constant delay less than half of the interval, *J. Math. Phys.*, **65(3)** (2024), No 032706.
- [27] F. Wang, C.-F. Yang, S. Buterin and N. Djurić, Inverse spectral problems for Dirac-type operators with global delay on a star graph, *Anal. Math. Phys.*, **14(2)** (2024), No 24.

Nebojša Djurić Faculty of Electrical Engineering, University of Banja Luka, Bosnia and Herzegovina.

E-mail: nebojsa.djuric@etf.unibl.org

Biljana Vojvodić Faculty of Mechanical Engineering, University of Banja Luka, Banja Luka, Bosnia and Herzegovina.

E-mail: biljana.vojvodic@mf.unibl.org