



Riemann solitons on Lorentzian generalized symmetric spaces

Mehdi Jafari and Shahroud Azami

Abstract. In this paper, we classify Riemann solitons on simply-connected 4-dimensional Lorentzian generalized symmetric spaces up to isometry. Then it is proved which of them is the gradient soliton. Also, we prove none of the potential vector fields of Riemann solitons are Killing vector fields.

Keywords. Conformal vector fields, generalized symmetric spaces, pseudo-Riemannian metrics.

1 Introduction

Hamilton [34] introduced the Ricci flow on a pseudo-Riemannian manifold (M, g) with the Ricci tensor S as follows

$$\frac{\partial}{\partial t} g = -2S. \quad (1.1)$$

Special solution to (1.1) is Ricci soliton [16] which is given by

$$\mathcal{L}_W g + S + \lambda g = 0, \quad (1.2)$$

for some constant λ and vector field W where $\mathcal{L}_W$ denotes the Lie derivative along W .

A Ricci soliton is a generalization of Einstein's metric and has applications in physics [1, 23, 24, 32, 38]. The pseudo-Riemannian geometry has more interesting properties than the Riemannian state and Ricci solitons have been studied on it [13, 17]. For example, in dimension three, there are some Riemannian homogeneous Ricci solitons [7, 41], but there are no left-invariant Riemannian Lie groups that admit Ricci solitons [31, 35, 42]. Also, there are several examples of the left-invariant three-dimensional Lorentzian Ricci solitons [13]. Many authors have generalized the Ricci flow and introduced new geometric flows. For instance, on manifold (M, g) with Riemann curvature tensor R the Riemann flow introduced by Udriște [46, 47] is as follows

$$\frac{\partial}{\partial t} G(t) = -2R(g(t)),$$

where $G = \frac{1}{2}g \odot g$ where for two $(0, 2)$ -tensors ω and θ product $\odot$ is defined by

$$\begin{aligned} (\omega \odot \theta)(Z_1, Z_2, Z_3, Z_4) &= \omega(Z_1, Z_4)\theta(Z_2, Z_3) + \omega(Z_2, Z_3)\theta(Z_1, Z_4) \\ &\quad - \omega(Z_1, Z_3)\theta(Z_2, Z_4) - \omega(Z_2, Z_4)\theta(Z_1, Z_3), \end{aligned}$$

for all vector fields Z_1, Z_2, Z_3, Z_4 . Manifold (M, g) is said to be a Riemann soliton (or RS) [36] and denoted by (M^n, g, μ, Y) if it admits

$$2R + \mu g \odot g + g \odot \mathcal{L}_Y g = 0, \quad (1.3)$$

for some vector field Y and constant μ . If $\mu > 0$ or $\mu < 0$ or $\mu = 0$ then the Riemann soliton is called expanding or shrinking or steady. A Riemann soliton is labeled as a gradient Riemann soliton if $Y = \text{grad}h$ for a smooth map h and the RS becomes

$$2R + \mu g \odot g + 2g \odot \nabla^2 h = 0.$$

When μ is a smooth function then RS and gradient RS is said to be an almost RS and an almost gradient RS, respectively. The RS is corresponding to the Riemann flow as a fixed point. From (1.3), we get

$$\begin{aligned} 2R(Z_1, Z_2, Z_3, Z_4) &= -2\mu [g(Z_1, Z_4)g(Z_2, Z_3) - g(Z_1, Z_3)g(Z_2, Z_4)] \\ &\quad - [g(Z_1, Z_4)\mathcal{L}_V g(Z_2, Z_3) + g(Z_2, Z_3)\mathcal{L}_V g(Z_1, Z_4)] \\ &\quad + [g(Z_1, Z_3)\mathcal{L}_V g(Z_2, Z_4) + g(Z_2, Z_4)\mathcal{L}_V g(Z_1, Z_3)]. \end{aligned} \quad (1.4)$$

Contracting Z_1 and Z_4 in the last equation, we have

$$2S(Z_2, Z_3) = -2((n-1)\mu + \text{div}V)g(Z_2, Z_3) - (n-2)\mathcal{L}_V g(Z_2, Z_3). \quad (1.5)$$

A lot of studies have been done on RSs on manifolds. For instance, Biswas et al. [10] investigated RS on a almost co-Kähler 3-dimensional manifold, Venkatesha et al. [30, 48] studied RS on contact geometry and almost Kenmotsu manifolds, and K. De and U. C. De [25] studied almost RS on para-Sasakian manifolds. For the Ricci solitons on the Walker manifolds we refer the reader to [14, 15]. Also, see [8, 11, 26].

We use the abbreviation GSS to represent generalized symmetric spaces, and FGSS for four-dimensional GSS. In a paper by Cerny and Kowalski [22], pseudo-Riemannian FGSS were classified into four classes denoted as A, B, C , and D . With the exception of type C , which is Lorentzian, the associated pseudo-Riemannian metrics have signatures of $(4, 0)$, $(2, 2)$, and $(0, 4)$. The geometric properties of these spaces, such as the Levi-Civita connection, curvature tensor, and Ricci tensor, were calculated in [20, 29]. Batat and Onda [9] conducted a study on algebraic Ricci solitons on pseudo-Riemannian FGSS. Numerous research works have been carried out on the geometric structures of GGSSs, including studies on homogeneous geodesics [28], curvature properties [20], harmonicity properties of invariant vector fields [19], Ricci solitons [12], and Ricci bi-conformal vector fields [2]. Also, see [33, 37, 39, 40, 44, 45, 18].

Motivated by mentioned works, we study the RSs on Lorentzian FGSSs up to isometry. The article is arranged as follows. We recall some essential notions about simply-connected non-symmetric Lorentzian FGSSs which be used throughout this paper in Section 2. In the Section 3, the main results and their proof are presented.

2 Preliminaries

Let (M, g) be a pseudo-Riemannian connected manifold and $x \in M$. If an isometry s_x of M is such that x is fixed isolated point, then it is called symmetry at the point x .

A family of isometries denoted by $\{s_x|x \in M\}$ is termed a regular s structure if it satisfies the condition $s_x \circ s_y = s_{s_x(y)} \circ s_x$ for all $x, y \in M$, where s_x represents a symmetry for all $x \in M$, and the mapping $M \times M \rightarrow M$, given by $(x, y) \mapsto s_x(y)$, is smooth. (M, g) that possesses at least one regular s structure is referred to as a GSS. In the context of GSS, if (M, g) is a GSS, it is homogeneous, and can be identified with $(G/H, g)$, where G is a subgroup of the isometry group $I(M)$ that acts transitively on M , and H denotes the isotropy group at a fixed point $x \in M$. A FGSS of type C is the underlying homogeneous space G/H , where

$$G = \begin{pmatrix} e^{-w} & 0 & 0 & x \\ 0 & e^w & 0 & y \\ 0 & 0 & 1 & z \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

A FGSS of type C is the space $\mathbb{R}^4(x, y, z, w)$ with the metric

$$g = \epsilon(e^{-2w}dx^2 + e^{2w}dy^2) + dzdw, \quad (2.1)$$

where $\epsilon = \pm 1$. The possible signatures are $(1, 3), (3, 1)$.

Now, let $(M = G/H, g)$ be a non-symmetric simply-connected FGSS of type C with coordinates (x, y, z, w) . Let ∇ be the Levi-Civita connection of (M, g) and R be its curvature tensor. The Ricci tensor is defined by

$$S(X_1, X_2) = \text{trace}(X_3 \rightarrow R(X_1, X_3)X_2). \quad (2.2)$$

Let

$$\partial_1 = \frac{\partial}{\partial x}, \quad \partial_2 = \frac{\partial}{\partial y}, \quad \partial_3 = \frac{\partial}{\partial z}, \quad \partial_4 = \frac{\partial}{\partial w}$$

be the coordinates vector fields on M . From [12], ∇ of M with respect to $\{\partial_i\}_{i=1}^4$ is given by

$$\nabla_{\partial_i}\partial_j = \begin{pmatrix} 2\epsilon e^{-2w}\partial_3 & 0 & 0 & -\partial_1 \\ 0 & -2\epsilon e^{2w}\partial_3 & 0 & \partial_2 \\ 0 & 0 & 0 & 0 \\ -\partial_1 & \partial_2 & 0 & 0 \end{pmatrix}, \quad (2.3)$$

and

$$\begin{aligned} R(\partial_2, \partial_4)\partial_4 &= \partial_2, & R(\partial_4, \partial_2)\partial_2 &= 2\epsilon e^{2w}\partial_3, \\ R(\partial_4, \partial_1)\partial_4 &= -\partial_1, & R(\partial_1, \partial_4)\partial_1 &= -2\epsilon e^{-2w}\partial_3. \end{aligned}$$

Hence, the Riemann curvature tensor is characterized by non-zero elements

$$\begin{aligned} R_{1414} &= -R_{1441} = -R_{4114} = R_{4141} = -\epsilon e^{-2w}, \\ R_{2424} &= -R_{2442} = -R_{4224} = R_{4242} = -\epsilon e^{2w} \end{aligned} \quad (2.4)$$

and the Ricci tensor is given by

$$S = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -2 \end{pmatrix}.$$

Along an arbitrary vector field $X = X_i \partial_i$ on manifold where $X_i = X_i(x, y, z, w)$, $i = 1, 2, 3, 4$ are smooth maps, we have

$$\begin{cases} (\mathcal{L}_X g)_{11} = 2\epsilon e^{-2w}(\partial_1 X_1 - X_4), & (\mathcal{L}_X g)_{12} = \epsilon(e^{2w}\partial_1 X_2 + e^{-2w}\partial_2 X_1), \\ (\mathcal{L}_X g)_{13} = \frac{1}{2}\partial_1 X_4 + \epsilon e^{-2w}\partial_3 X_1, & (\mathcal{L}_X g)_{14} = \frac{1}{2}\partial_1 X_3 + \epsilon e^{-2w}\partial_4 X_1, \\ (\mathcal{L}_X g)_{22} = 2\epsilon e^{2w}(X_4 + \partial_2 X_2), & (\mathcal{L}_X g)_{23} = \frac{1}{2}\partial_2 X_4 + \epsilon e^{2w}\partial_3 X_2, \\ (\mathcal{L}_X g)_{24} = \frac{1}{2}\partial_2 X_3 + \epsilon e^{2w}\partial_4 X_2, & (\mathcal{L}_X g)_{33} = \partial_3 X_4, \\ (\mathcal{L}_X g)_{34} = \frac{1}{2}(\partial_3 X_3 + \partial_4 X_4), & (\mathcal{L}_X g)_{44} = \partial_4 X_3. \end{cases} \quad (2.5)$$

3 RS on FGSS of type C

Now, we classify all RSs on non-symmetric simply-connected FGSSs of type C . Using (1.4), (M, g, μ, X) is a Riemann soliton if and only if

$$\begin{aligned} 2R_{1414} &= g_{11}(\mathcal{L}_X g)_{44}, \\ 2R_{2424} &= g_{22}(\mathcal{L}_X g)_{44}, \\ 2R_{3434} &= -2\mu g_{34}^2 - 2g_{34}(\mathcal{L}_X g)_{34}, \\ 2R_{1212} &= 2\mu g_{11}g_{22} + g_{11}(\mathcal{L}_X g)_{22} + g_{22}(\mathcal{L}_X g)_{11}, \\ 2R_{1313} &= g_{11}(\mathcal{L}_X g)_{33}, \\ 2R_{1214} &= g_{11}(\mathcal{L}_X g)_{24}, \\ 2R_{1213} &= g_{11}(\mathcal{L}_X g)_{23}, \\ 2R_{1223} &= -g_{22}(\mathcal{L}_X g)_{13}, \\ 2R_{1224} &= -g_{22}(\mathcal{L}_X g)_{14}, \\ 2R_{1314} &= 2\mu g_{11}g_{34} + g_{11}(\mathcal{L}_X g)_{34} + g_{34}(\mathcal{L}_X g)_{11}, \\ 2R_{1324} &= g_{34}(\mathcal{L}_X g)_{12}, \\ 2R_{1334} &= g_{34}(\mathcal{L}_X g)_{13}. \end{aligned} \quad (3.1)$$

Applying (2.1), (2.4), and (2.5) in the equations (3.1), we have

$$\partial_4 X_3 = -2, \quad (3.2)$$

$$\frac{1}{2}(\partial_3 X_3 + \partial_4 X_4) = -\mu, \quad (3.3)$$

$$\mu + \partial_2 X_2 + \partial_1 X_1 = 0, \quad (3.4)$$

$$\partial_3 X_4 = 0, \quad (3.5)$$

$$\frac{1}{2}\partial_2 X_3 + \epsilon e^{2w}\partial_4 X_2 = 0, \quad (3.6)$$

$$\frac{1}{2}\partial_2 X_4 + \epsilon e^{2w}\partial_3 X_2 = 0, \quad (3.7)$$

$$\frac{1}{2}\partial_1 X_4 + \epsilon e^{-2w}\partial_3 X_1 = 0, \quad (3.8)$$

$$\frac{1}{2}\partial_1 X_3 + \epsilon e^{-2w}\partial_4 X_1 = 0, \quad (3.9)$$

$$2\mu + \frac{1}{2}(\partial_3 X_3 + \partial_4 X_4) + 2(\partial_1 X_1 - X_4) = 0, \quad (3.10)$$

$$e^{2w}\partial_1 X_2 + e^{-2w}\partial_2 X_1 = 0. \quad (3.11)$$

Equations (3.2) and (3.5) imply that

$$X_3 = -2w + G(x, y, z), \quad X_4 = F(x, y, w), \quad (3.12)$$

for smooth maps F and G . Deriving equations (3.6) and (3.9) with respect to w , one gets

$$\partial_{44}^2 X_1 - 2\partial_4 X_1 = 0, \quad (3.13)$$

$$\partial_{44}^2 X_2 + 2\partial_4 X_2 = 0. \quad (3.14)$$

Solving these equations and using equations (3.6) and (3.9), we deduce

$$X_1 = H(x, y, z) - \frac{1}{4\epsilon} e^{2w} \partial_1 G(x, y, z), \quad (3.15)$$

$$X_2 = K(x, y, z) + \frac{1}{4\epsilon} e^{-2w} \partial_2 G(x, y, z) \quad (3.16)$$

for some smooth functions H and K . Utilizing (3.15) and (3.16) in (3.11), we conclude $e^{2w} \partial_1 K(x, y, z) + e^{-2w} \partial_2 H(x, y, z) = 0$. Hence $\partial_1 K(x, y, z) = 0$ and $\partial_2 H(x, y, z) = 0$. Therefore,

$$X_1 = L(x, z) - \frac{1}{4\epsilon} e^{2w} \partial_1 G(x, y, z), \quad (3.17)$$

$$X_2 = I(y, z) + \frac{1}{4\epsilon} e^{-2w} \partial_2 G(x, y, z) \quad (3.18)$$

for some smooth functions L and I . Deriving (3.7) and (3.8) with respect to z , it follows $\partial_{33}^2 X_1 = 0$ and $\partial_{33}^2 X_2 = 0$. It follows that

$$L(x, y) = L_1(x)z + L_2(x), \quad \partial_1 G(x, y, z) = G_1(x, y)z + G_2(x, y), \quad (3.19)$$

$$I(x, y) = I_1(y)z + I_2(y), \quad \partial_2 G(x, y, z) = G_3(x, y)z + G_4(x, y), \quad (3.20)$$

and

$$X_1 = L_1(x)z + L_2(x) - \frac{1}{4\epsilon} e^{2w} (G_1(x, y)z + G_2(x, y)), \quad (3.21)$$

$$X_2 = I_1(y)z + I_2(y) + \frac{1}{4\epsilon} e^{-2w} (G_3(x, y)z + G_4(x, y)). \quad (3.22)$$

Equations (3.19) and (3.20) yield

$$\partial_2 G_1(x, y) = \partial_1 G_3(x, y), \quad \partial_2 G_2(x, y) = \partial_1 G_4(x, y). \quad (3.23)$$

Applying (3.21) and (3.22) in (3.7) and (3.8), we infer

$$\frac{1}{2} \partial_1 F(x, y, w) + \epsilon e^{-2w} \left(L_1(x) - \frac{1}{4\epsilon} e^{2w} G_1(x, y) \right) = 0, \quad (3.24)$$

$$\frac{1}{2} \partial_2 F(x, y, w) + \epsilon e^{2w} \left(I_1(y) + \frac{1}{4\epsilon} e^{-2w} G_3(x, y) \right) = 0. \quad (3.25)$$

Deriving (3.24) and (3.25) with respect to y and x , respectively, we find

$$\frac{1}{2} \partial_{12}^2 F(x, y, w) - \frac{1}{4} \partial_2 G_1(x, y) = 0, \quad (3.26)$$

$$\frac{1}{2} \partial_{21}^2 F(x, y, w) + \frac{1}{4} \partial_1 G_3(x, y) = 0. \quad (3.27)$$

Equations (3.26) and (3.27) lead to $-\partial_2 G_1(x, y) = \partial_1 G_3(x, y)$. Then (3.23) leads to $\partial_2 G_1(x, y) = \partial_1 G_3(x, y) = 0$ and $G_1(x, y) = G_5(x)$ and $G_3(x, y) = G_6(y)$ for some smooth functions G_5 and G_6 . Using (3.3), (3.4), and (3.10), we have

$$\partial_1 X_1 - \partial_2 X_2 - 2X_4 = 0. \quad (3.28)$$

Upon differentiating equation (3.28) with respect to z it follows

$$\partial_{13}^2 X_1 - \partial_{23}^2 X_2 = 0. \quad (3.29)$$

Substituting (3.21) and (3.22) in (3.29), we get

$$L_1'(x) - \frac{1}{4\epsilon} e^{2w} G_5'(x) - I_1'(y) - \frac{1}{4\epsilon} e^{-2w} G_6'(y) = 0. \quad (3.30)$$

The last equation yields $L_1'(x) = I_1'(y)$ and $G_5'(x) = G_6'(y) = 0$, then

$$L_1(x) = a_1 x + a_2, \quad I_1(y) = a_1 y + a_3, \quad G_5(x) = a_4, \quad G_6(y) = a_5 \quad (3.31)$$

for some constants a_1, a_2, a_3, a_4 and a_5 . Upon differentiating equation (3.28) regarding to x it follows that

$$L_2''(x) - \frac{1}{4\epsilon} e^{2w} \partial_{11}^2 G_2(x, y) - \frac{1}{4\epsilon} e^{-2w} \partial_{12}^2 G_4(x, y) + 4\epsilon e^{-2w} (a_1 x + a_2) - a_4 = 0. \quad (3.32)$$

Equation (3.32) leads to

$$\partial_{11}^2 G_2(x, y) = 0, \quad \partial_{12}^2 G_4(x, y) = 16(a_1 x + a_2), \quad L_2''(x) = a_4. \quad (3.33)$$

Upon differentiating equation (3.28) regarding to y , we arrive at

$$-I_2''(y) - \frac{1}{4\epsilon} e^{2w} \partial_{12}^2 G_2(x, y) - \frac{1}{4\epsilon} e^{-2w} \partial_{22}^2 G_4(x, y) + 4\epsilon e^{2w} (a_1 y + a_3) + a_5 = 0. \quad (3.34)$$

Equation (3.34) leads to

$$\partial_{12}^2 G_2(x, y) = 16(a_1 y + a_3), \quad \partial_{22}^2 G_4(x, y) = 0, \quad I_2''(y) = a_5. \quad (3.35)$$

From (3.23), (3.33), and (3.35), we conclude $a_1 = a_2 = a_3 = 0$

$$\begin{aligned} G_2(x, y) &= 16\left(\frac{1}{2}a_1 y^2 + a_3 y + a_6\right)x + 8a_2 y^2 - a_7 y + a_8, \\ G_4(x, y) &= 16\left(\frac{1}{2}a_1 x^2 + a_2 x + a_9\right)y + 8a_3 x^2 - a_7 x + a_{10}, \\ L_2(x) &= \frac{1}{2}a_4 x^2 + a_{11}x + a_{12}, \quad I_2(y) = \frac{1}{2}a_5 y^2 + a_{13}y + a_{14}, \end{aligned}$$

for some constants $a_6, \dots, a_{14}$. Also, from (3.24) and (3.25), we deduce

$$F(x, y, w) = -2\epsilon e^{-2w} \left(\frac{1}{2}a_1 x^2 + a_2 x\right) + \frac{1}{2}a_4 x - 2\epsilon e^{2w} \left(\frac{1}{2}a_1 y^2 + a_3 y\right) - \frac{1}{2}a_5 y + F_1(w) \quad (3.36)$$

for some smooth function F_1 . Applying (3.28), it follows that

$$F_1(w) = \frac{1}{2} (a_{11} - a_{13} - 4\epsilon a_6 e^{2w} - 4\epsilon a_9 e^{-2w}). \quad (3.37)$$

Equation (3.3) implies that

$$\mu = -\frac{1}{2} \left(\partial_3 G(x, y, z) + 4\epsilon e^{-2w} \left(\frac{1}{2} a_1 x^2 + a_2 x + a_9 \right) - 4\epsilon e^{2w} \left(\frac{1}{2} a_1 y^2 + a_3 y + a_6 \right) \right). \quad (3.38)$$

Equation (3.4) implies that

$$\begin{aligned} \mu = & -(2a_1 z + a_4 x + a_{11} + a_5 y + a_{13} + 4\epsilon e^{-2w} \left(\frac{1}{2} a_1 x^2 + a_2 x + a_9 \right) \\ & - 4\epsilon e^{2w} \left(\frac{1}{2} a_1 y^2 + a_3 y + a_6 \right)). \end{aligned} \quad (3.39)$$

Compare (3.38) and (3.39) gives $a_1 = a_2 = a_3 = a_6 = a_9 = 0$ and $\partial_3 G(x, y, z) = 2(a_4 x + a_{11} + a_5 y + a_{13})$. Applying (3.39) in (3.10) we get $a_{11} = 0$. Using $\partial_3 G(x, y, z) = 2(a_4 x + a_5 y + a_{13})$, (3.19), and (3.20), we obtain $a_4 = a_5 = 0$,

$$G(x, y, z) = -a_7 xy + a_8 x + a_{10} y + 2a_{13} z + a_{14} \quad (3.40)$$

for some constant a_{14} .

Hence, we have the next consequence:

Theorem 3.1. *A FGSS of type C admit almost Riemann soliton (M, g, μ, X) with $X = X_i \partial_i$ if and only if $\mu = b_1$ and*

$$\begin{cases} X_1 = b_2 - \frac{1}{4\epsilon} e^{2w} (b_3 y + b_4), \\ X_2 = -b_1 y + b_5 + \frac{1}{4\epsilon} e^{-2w} (b_3 x + b_6), \\ X_3 = -2w + b_3 xy + b_4 x + b_6 y - 2b_1 z + b_7, \\ X_4 = \frac{1}{2} b_1, \end{cases} \quad (3.41)$$

where $b_1, b_2, \dots, b_7 \in \mathbb{R}$.

Theorem (3.1) gives:

Corollary 3.2. *Any almost Riemann soliton on FGSS of type C is a Riemann soliton.*

Now, we investigate which of RSs are gradient RS. For this end, we consider the potential vector field of a gradient RS is $X = \nabla f$ for some smooth function f on a FGSS of type C. On a FGSS, we have

$$\nabla f = \epsilon e^{2w} (\partial_1 f) \partial_1 + \epsilon e^{-2w} (\partial_2 f) \partial_2 + 2(\partial_4 f) \partial_3 + 2(\partial_3 f) \partial_4. \quad (3.42)$$

From (3.41) and (3.42), we have

$$\partial_1 f = \epsilon b_2 e^{-2w} - \frac{1}{4} (b_3 y + b_4), \quad (3.43)$$

$$\partial_2 f = \epsilon e^{2w} (-b_1 y + b_5) + \frac{1}{4} (b_3 x + b_6), \quad (3.44)$$

$$\partial_3 f = \frac{1}{4} b_1, \quad (3.45)$$

$$\partial_4 f = \frac{1}{2} (-2w + b_3 xy + b_4 x + b_6 y - 2b_1 z + b_7). \quad (3.46)$$

By taking derivative of the equations (3.43) and (3.46) with respect to w and x , respectively, we obtain $-2\epsilon b_2 e^{-2w} = \frac{1}{2}(b_3 y + b_4)$ which implies that $b_2 = b_3 = b_4 = 0$. Also, taking derivative of (3.44) and (3.46) with respect to w and y , respectively, we deduce $\epsilon e^{2w}(-b_1 y + b_5) = \frac{1}{2}b_6$ which yields to $b_1 = b_5 = b_6 = 0$. Thus

$$\partial_1 f = \partial_2 f = \partial_3 f = 0, \text{ and } \partial_4 f = \frac{b_7}{2}, \quad (3.47)$$

and

$$f = \frac{b_7}{2}w + b_8, \quad (3.48)$$

where $b_8 \in \mathbb{R}$. Therefore, we have:

Corollary 3.3. *A FGSS of type C admits gradient RS $(M, g, \mu, \nabla f)$ if and only if $f = \frac{a}{2}w + b$, where $a, b \in \mathbb{R}$.*

Remark 1. A Killing vector field on (M, g) is a vector field X such that

$$\mathcal{L}_X g = 0.$$

Hence, from Theorem 3.1 we deduce that any potential vector fields of almost Riemann soliton on FGSS of type C is not Killing vector field because $(\mathcal{L}_X g)_{44} = \partial_4 X_3 = -2 \neq 0$. Also, is not conformal vector field.

Remark 2. A Ricci collineation vector field [21] is the vector field X such that $\mathcal{L}_X S = 0$. The Lie-derivative of S regarding to $X = X_i \partial_i$ on FGSS of type C is represented by

$$\mathcal{L}_X S = \begin{pmatrix} 0 & 0 & 0 & -2\partial_1 X_4 \\ 0 & 0 & 0 & -2\partial_2 X_4 \\ 0 & 0 & 0 & -2\partial_3 X_4 \\ -2\partial_1 X_4 & -2\partial_2 X_4 & -2\partial_3 X_4 & -4\partial_4 X_4 \end{pmatrix}. \quad (3.49)$$

Using Theorem 3.1, the potential vector fields of Riemann solitons on FGSS of type C become Ricci collineation vector field.

Remark 3. Ricci bi-conformal vector field [27] on a pseudo-Riemannian manifold (M, g) is a vector field X such that

$$\mathcal{L}_X g = \alpha g + \beta S, \quad \mathcal{L}_X S = \alpha S + \beta g, \quad (3.50)$$

for smooth maps β and α . Also, see [2, 3, 4, 5, 6, 43]. Using Theorem 3.1, $\mathcal{L}_X S = 0$ which leads to $\alpha = \beta = 0$. But $\mathcal{L}_X g \neq 0$. Therefore, the potential vector fields of Riemann solitons on FGSSs of type C are not Ricci bi-conformal vector fields.

Remark 4. If a vector field X will be a potential vector field of a Ricci soliton on FGSSs of type C then the equation (1.2) implies that it satisfies in the following system.

$$\begin{cases} 2\epsilon e^{-2w}(\partial_1 X_1 - X_4) = \epsilon \lambda e^{-2w}, & \epsilon(e^{2w} \partial_1 X_2 + e^{-2w} \partial_2 X_1) = 0, \\ \frac{1}{2} \partial_1 X_4 + \epsilon e^{-2w} \partial_3 X_1 = 0, & \frac{1}{2} \partial_1 X_3 + \epsilon e^{-2w} \partial_4 X_1 = 0, \\ 2\epsilon e^{2w}(X_4 + \partial_2 X_2) = \epsilon \lambda e^{2w}, & \frac{1}{2} \partial_2 X_4 + \epsilon e^{2w} \partial_3 X_2 = 0, \\ \frac{1}{2} \partial_2 X_3 + \epsilon e^{2w} \partial_4 X_2 = 0, & \partial_3 X_4 = 0, \\ \frac{1}{2}(\partial_3 X_3 + \partial_4 X_4) = \lambda, & \partial_4 X_3 = 2. \end{cases} \quad (3.51)$$

Applying Theorem 3.1, we conclude that any potential vector fields of almost Riemann soliton on FGSS of type C is not potential vector fields of Ricci soliton because $\partial_4 X_3 = -2 \neq 2$.

References

- [1] M. M. Akbar and E. Woolgar, Ricci solitons and Einstein-Scalar field theory, *Classical Quantum Gravity*, **26**(5) (2009), 055015, 14pp.
- [2] S. Azami, U. C. De, Ricci bi-conformal vector fields on Lorentzian four-dimensional generalized symmetric spaces, *Filomat*, **38**(4) (2024), 3157-3164.
- [3] S. Azami, and U. C. De, Ricci bi-conformal vector fields on Lorentzian five-dimensional two-step nilpotent Lie groups, *Hacet. J. Math. Stat.* **53**(4) (2024), 1118-1129
- [4] S. Azami, and G. Fasihi-Ramandi, Ricci bi-conformal vector fields on Siklos spacetimes, *Mathematics. Interdisciplinary research*, **9**(1) (2024), 45-76.
- [5] S. Azami, and M. Jafari, Ricci bi-conformal vector fields on homogeneous Godel-type spacetimes, *J. nonlinear Math. Phys.* **30** (2023), 1700-1718.
- [6] S. Azami and M. Jafari, Ricci solitons and Ricci bi-conformal vector fields on the Lie group $\mathbb{H}^2 \times \mathbb{R}$, *Report on mathematical physics*, **93**(2) (2024), 231-239.
- [7] P. Baird and L. Danielo, Three-dimensional Ricci solitons which project to surfaces, *J. Reine Angew. Math.*, **2007**(608) (2007), 65-91.
- [8] M.R. Bakshi, K. K. Baishya, Four classes of Riemann solitons on α -cosymplectic manifolds, *Afr. Mat.*, **32** (2020), 577-588.
- [9] W. Batat and K. Onda, Four-dimensional pseudo-Riemannian generalized symmetric spaces which are algebraic Ricci solitons, *Results Math.*, **64** (2013), 253-267.
- [10] G.G. Biswas, X. Chen, U. C. De, Riemann solitons on almost co-Kähler manifolds, *Filomat*, **36**(4) (2022), 1403-1413.
- [11] A.M. Blaga, Remarks on almost Riemann solitons with gradient or torse-forming vector field. *Bulletin of the Malaysian Mathematical Sciences Society*, **44** (2020), 3215 - 3227.
- [12] A. Bouharis and B. Djebbar, Ricci solitons on Lorentzian four-dimensional generalized symmetric spaces, *Journal of mathematical physics, analysis, geometry*, **14**(2) (2018), 132-140.
- [13] M. Brozos-Vázquez, G. Calvaruso, E. García-Río and S. Gavino-Fernández, Three-dimensional Lorentzian homogeneous Ricci solitons, *Israel J. Math.*, **188** (2012), 385-403.
- [14] M. Brozos-Vázquez, E. Garcia-Río, and S. Gavino-Fernandez, Locally conformally flat Lorentzian gradient Ricci solitons, *J. Geom. Anal.* **23** (2013), 1196-1212.
- [15] M. Brozos-Vázquez, E. Garcia-Río, P. Gilkey, S. Nikčević, and R. Vázquez-Lorenzo, *The Geometry of Walker Manifolds*, Vol. 5 of *Synthesis Lectures on Mathematics and Statics*, Morgan and Claypool, San Rafael, CA, 2009.
- [16] C. Calin and M. Crasmăreanu, From the Eisenhart problem to Ricci solitons in f -Kenmotsu manifolds, *Bull. Malays. Math. Soc.* **33**(3) (2010), 361-368.
- [17] G. Calvaruso, Homogeneous structures on three-dimensional homogeneous Lorentzian manifolds, *J. Geom. Phys.*, **57** (2007), 1279-1291.

- [18] G. Calvaruso, Symplectic, complex and Kähler structures on four-dimensional generalized symmetric spaces, *Differential geometry and its applications*, **29** (2011), 758-769.
- [19] G. Calvaruso, Harmonicity of invariant vector fields on four-dimensional generalized symmetric spaces, *Center. Eur. J. Math.*, **10** (2012), 411-425.
- [20] G. Calvaruso and B. De Leo, Curvature properties of four-dimensional generalized symmetric spaces, *J. Geom.*, **90** (2008), 30-46.
- [21] U. Camci, H. Baysal, I. Tarhan, I. Yilmaz and I. Yavuz, Ricci collineations of the Bianchi type I and III, and Kantowski-Sachs spacetimes, *Int. J. Modern Physics D*, **10**(5) (2001), 751-765.
- [22] J. Cerny and O. Kowalski, Classification of generalized symmetric pseudo-Riemannian spaces of dimension $n = 4$, *Tensor N.S.* **38** (1982), 256-267.
- [23] T. Chave and G. Valent, Quasi-Einstein metrics and their renormalizability properties, *Helv. Phys. Acta.* **69** (1996), 344-347.
- [24] T. Chave, G. Valent, On a class of compact and non-compact quasi-Einstein metrics and their renormalizability properties, *Nuclear Phys. B.* **478** (1996) 758-778.
- [25] K. De and U. C. De, Riemann solitons on para-Sasakian geometry, *Carpathian Mathematical Publications*, **14**(2) (2022), 395-405.
- [26] K. De, U.C. De, A note on almost Riemann Solitons and gradient almost Riemann Solitons, *Afr. Mat.* **33**(74) (2022). <https://doi.org/10.1007/s13370-022-01010-y>.
- [27] U. C. De, A. Sardar, and A. Sarkar, Some conformal vector fields and conformal Ricci solitons on $N(k)$ -contact metric manifolds, *AUT J. Math. Com.*, **2**(1) (2021), 61-71.
- [28] B. De Leo and R. A. Marinosci, Homogeneous geodesics of four-dimensional generalized symmetric pseudo-Riemannian spaces, *Publ. Math. Debrecen.* **73** (2008), 341-360.
- [29] B. De Leo and J. V. Der Veken, Totally geodesic hypersurfaces of four-dimensional generalized symmetric spaces, *Geometriae Dedicata*, **159**(1) (2012), 373-387.
- [30] M. N. Devaraja, H. A. Kumara and V. Venkatesha, Riemannian soliton within the framework of contact geometry, *Quaestiones Mathematicae*, **44** (2021), 637-651.
- [31] L. F. di Cerbo, Generic properties of homogeneous Ricci soliton, *Advances in Geometry*, **14**(2) (2014), 225-237.
- [32] D. H. Friedan, Nonlinear models in $2 + \epsilon$ dimensions, *Ann. Physics*, **163**(2) (1985), 318-419.
- [33] A. Gray, Riemannian manifolds with geodesic symmetries of order 3, *J. Differential Geom.*, **7** (1970), 343-369.
- [34] R. S. Hamilton, The Ricci flow on surfaces, *Mathematics and general relativity, Contemporary Mathematics*, **71** (1988), 237-261.
- [35] S. Hervik, Ricci nilsoliton black holes, *J. Geom. Phys.*, **58**(9) (2008), 1253-1264.
- [36] I. E. Hirićă and C. Udriște, Ricci and Riemann solitons, *Balkan J. Geom. Appl.* **21**(2) (2016), 35-44.

- [37] V. G. Kac, Graded Lie algebras and symmetric spaces, Funk. Anal. Prilozhen, **2** (1968), 93-94.
- [38] A. L. Kholodenko, Towards physically motivated proofs of the Poincaré and geometrization conjectures, J. Geom. Phys., **58** (2) (2008), 259-290.
- [39] D. Kotschick and S. Terzić, On formality of generalized symmetric space, Math. Proc. Cambridge Philos. Soc., **134** (2003), 491-505.
- [40] O. Kowalski, Generalized symmetric spaces, in: Lectures Notes in Math. Springer-Verlag. Berlin, Heidelberg, NewYork, 1980.
- [41] J. Lauret, Ricci solitons solvmanifolds, Journal für die reine und angewandte Mathematik, **2011**(650) (2011), 1-21.
- [42] T. L. Payne, The existence of soliton metrics for nilpotent Lie groups, Geom. Dedicata, **145** (2010), 71-88.
- [43] M. Sohrabpour, S. Azami, Ricci bi-conformal vector fields on Lorentzian Walker manifolds of low dimension. Lobachevskii J Math., **44** (2023), 5437-5443. <https://doi.org/10.1134/S1995080223120338>.
- [44] S. Terzić, Real cohomology of generalized symmetric spaces, Fundam, Prikl Mat., **7** (2001), 131-157.
- [45] S. Terzić, Pontryagin classes of generalized symmetric spaces, Mat. Zametki, **69** (2001), 613-621, Translition in Math. Notes, **69** (2001), 559-566.
- [46] C. Udrişte, Riemann flow and Riemann wave, Ann. Univ. Vest Timisoara. Ser. Mat. Inf. **48**(1-2) (2010), 256-274.
- [47] C. Udrişte, Riemann flow and Riemann wave via bialternate product Riemannian metric, Preprint arXiv:1112.4279 [math. Ap] (2012).
- [48] V. Venkatesha, H. A. Kumara and M. N. Devaraja, Riemann solitons and almost Riemann solitons on almost Kenmotsu manifolds, International Journal of Geometric Methods in Modern Physics, **17** (2020), 2050105.

Mehdi Jafari Department of Mathematics, Payame Noor University, PO BOX 19395-4697, Tehran, Iran.

E-mail: m.jafarii@pnu.ac.ir

Shahroud Azami Department of pure Mathematics, Faculty of Science, Imam Khomeini International University, Qazvin, Iran.

E-mail: azami@sci.ikiu.ac.ir